
UK Renal Registry 16th Annual Report: Appendix I Acronyms and Abbreviations used in the Report

| | |
|-----------------|------------------------------------------------------------|
| ACE (inhibitor) | Angiotensin converting enzyme (inhibitor) |
| AKI | Acute kidney injury |
| ANZDATA | Australia and New Zealand Dialysis and Transplant Registry |
| APD | Automated peritoneal dialysis |
| ADPKD | Autosomal dominant polycystic kidney disease |
| APKD | Adult polycystic kidney disease |
| ATTOM | Access to transplant and transplant outcome measures |
| AV | Arteriovenous |
| AVF | Arteriovenous fistula |
| AVG | Arteriovenous graft |
| BAPN | British Association of Paediatric Nephrology |
| BCG | Bromocresol green |
| BCP | Bromocresol purple |
| BMD | Bone mineral disease |
| BMI | Body mass index |
| BP | Blood pressure |
| BSI | Blood stream infection |
| BTS | British Transplant Society |
| Ca | Calcium |
| CAB | Clinical Affairs Board (Renal Association) |
| CABG | Coronary artery bypass grafting |
| CAPD | Continuous ambulatory peritoneal dialysis |
| CCL | Clinical Computing Limited |
| CCPD | Cycling peritoneal dialysis |
| CDI | Clostridium difficile infection |
| Chol | Cholesterol |
| CHr | Target reticulocyte Hb content |
| CI | Confidence interval |
| CK | Creatine kinase |
| CKD | Chronic kidney disease |
| CKD-EPI | Chronic kidney disease epidemiology collaboration |
| CK-MB | Creatine kinase isoenzyme MB |
| COPD | Chronic obstructive pulmonary disease |
| CRF | Chronic renal failure |
| cRF | Calculated HLA antibody reaction frequency |
| CRP | C-reactive protein |
| CVVH | Continuous veno-venous haemofiltration |
| CXR | Chest x-ray |
| DBP | Diastolic blood pressure |
| DCCT | Diabetes Control and Complications Trial |
| DH | Department of Health |

| | |
|-----------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| DM | Diabetes mellitus |
| DOB | Date of birth |
| DOPPS | Dialysis Outcomes and Practice Patterns Study |
| E&W | England and Wales |
| E, W & NI | England, Wales and Northern Ireland |
| EBPG | European Best Practice Guidelines |
| ECG | Electrocardiogram |
| EDTA | European Dialysis and Transplant Association |
| EF | Error factor |
| eGFR | Estimated glomerular filtration rate |
| E_i | Expected cases in area i |
| ECD | Extended Criteria Donor |
| EDTA | European Dialysis and Transplant Association |
| eKt/V | Equilibrated Kt/V |
| EPO | Erythropoietin |
| ERA | European Renal Association |
| ERA-EDTA | European Renal Association-European Dialysis and Transplant Association |
| ERF | Established renal failure |
| ESA | Erythropoiesis stimulating agent |
| ESRD | End stage renal disease |
| ESRF | End stage renal failure |
| EWNI | England, Wales and Northern Ireland |
| Ferr | Ferritin |
| FEV1 | Forced expiratory volume in 1 second |
| FVC | Forced vital capacity |
| GFR | Glomerular filtration rate |
| GH | Growth hormone |
| GN | Glomerulonephritis |
| HA | Health Authority |
| HB | Health board |
| Hb | Haemoglobin |
| HbA1c | Glycated Haemoglobin |
| HBeAg | Hepatitis B e antigen |
| HCAI-DCS | Healthcare-associated infection data collection system |
| HD | Haemodialysis |
| HDF | Haemodialysis filtration |
| HDL | High-density lipoprotein |
| HLA | Human leucocyte antigen |
| HPA | Health Protection Agency |
| HQIP | Health Quality Improvement Partnership |
| HR | Hazard ratio |
| HRC | Hypochromic red blood cells |
| Ht | Height |
| ICU | Intensive care unit |
| IDMS | Isotope dilution mass spectrometry |
| IDOPPS | International Dialysis Outcomes and Practice Patterns Study |
| IFCC | International Federation of Clinical Chemistry & Laboratory Medicine |
| IHD | Ischaemic heart disease |
| IMD | Index of Multiple Deprivation |
| IOTF | International Obesity Taskforce |
| IPD | Intermittent peritoneal dialysis |
| IQR | Inter-quartile range |
| ISPD | International Society for Peritoneal Dialysis |
| IT | Information technology |
| IU | International units |
| IV | Intra venous |
| KDIGO | Kidney Disease: Improving Global Outcomes |
| KDOQI | Kidney Disease Outcomes Quality Initiative |
| KM | Kaplan Meier |
| Kt/V | Ratio between the product of urea clearance (K, in ml/min) and dialysis session duration (t, in minutes) divided by the volume of distribution of urea in the body (V, in ml) |

| | |
|----------------|--------------------------------------------------------------------------------|
| LA | Local Authority |
| LCL | Lower confidence limit |
| LDL | Low-density lipoprotein |
| LTFU | Lost to follow-up |
| M:F | Male:Female |
| MAP | Mean arterial blood pressure |
| MDRD | Modification of diet in renal disease |
| MI | Myocardial infarction |
| MMF | Mycophenolate mofetil |
| MRSA | Methicillin resistant Staphylococcal aureus |
| MSSA | Methicillin sensitive Staphylococcal aureus |
| N | Number |
| N Ireland | Northern Ireland |
| NE | North East |
| NEQAS | UK National External Quality Assessment Scheme |
| NHBPEP | National high blood pressure education programme |
| NHS | National Health Service |
| NHS BT | National Health Service Blood and Transplant |
| NI | Northern Ireland |
| NICE | National Institute for Health and Clinical Excellence |
| NISRA | Northern Ireland Statistic and Research Agency |
| NMO | Non-mixed origin |
| NRS | National Records of Scotland |
| NSF | National service framework |
| NTC | Non-tunnelled dialysis catheter |
| NTL | Non-tunnelled line |
| NW | North West |
| O/E | Observed/expected |
| ODT | Organ Donation and Transplantation (a Directorate of NHS Blood and transplant) |
| O _i | Observed cases in area i |
| ONS | Office for National Statistics |
| OR | Odds ratio |
| PAS | Patient Administration System |
| PCT | Primary Care Trust |
| PD | Peritoneal dialysis |
| PHE | Public Health England |
| Phos | Phosphate |
| PIAG | Patient Information Advisory Group |
| PKD | Polycystic kidney disease |
| PMARP | Per million age related population |
| PMCP | Per million child population |
| PMP | Per million population |
| PP | Pulse pressure |
| PRD | Primary renal disease |
| PTH | Parathyroid hormone |
| PUV | Posterior urethral valves |
| PVD | Peripheral vascular disease |
| QOF | Quality and Outcomes Framework |
| QUEST | Quality European Studies |
| RA | Renal Association |
| rhGH | Recombinant human growth hormone |
| RI | Royal Infirmary |
| RNSF | Renal National Service Framework (or NSF) |
| RR | Relative risk |
| RRDSS | Renal Registry data set specification |
| RRT | Renal replacement therapy |
| RVD | Renovascular disease |
| SAR | Standardised acceptance ratio (=O/E) |
| SAS | Statistical Analysis System |
| SBP | Systolic blood pressure |
| SD | Standard deviation |

| | |
|--------|------------------------------------------------------|
| SES | Socio-economic status |
| SHA | Strategic health authority |
| SHARP | Study of Heart and Renal Protection |
| SI | System International (units) |
| SMR | Standardised mortality ratios |
| spKt/V | Single pool Kt/V |
| SPR | Standardised prevalence ratio (=O/E) |
| SR | Standardised ratio (used to cover either SAR or SPR) |
| SUS | Secondary uses service |
| SW | South West |
| TC | Tunnelled dialysis catheter |
| TL | Tunnelled line |
| TSAT | Transferrin saturation |
| TWL | Transplant waiting list |
| Tx | Transplant |
| UCL | Upper confidence limit |
| UK | United Kingdom |
| UKRR | UK Renal Registry |
| UKT | UK Transplant (now ODT) |
| URR | Urea reduction ratio |
| US | United States |
| USA | United States of America |
| USRDS | United States Renal Data System |
| WHO | World health organization |
| Wt | Weight |
