
**Commentary on NICE guideline (NG136) ‘Hypertension in adults:
diagnosis and management’
including proposals for blood pressure management in patients
with chronic kidney disease**

Final version: March 2021
Review date: March 2026

Authors:

Dr Richard Fish

University Hospitals of North Midlands NHS Trust

Dr Nihil Chitalia

Dartford and Gravesham NHS Trust

Dr Timothy Doulton

East Kent Hospitals University NHS Foundation Trust

Ms Katie Durman

Barts Health NHS Trust

(on behalf of the British Dietetic Association, Renal Nutrition Group)

Ms Elizabeth Lamerton

Salford Royal Foundation Trust

(on behalf of the UK Renal Pharmacy Group)

Dr Eirini Lioudaki

King's College Hospital, London

Dr Adam MacDiarmid-Gordon

Brighton and Sussex University Hospitals NHS Trust

Professor Patrick Mark

Queen Elizabeth University Hospital, Glasgow

Dr Laura Ratcliffe

East and North Hertfordshire NHS Trust

Dr Charlie Tomson

Freeman Hospital, Newcastle upon Tyne

Professor Indranil Dasgupta

University Hospitals Birmingham NHS Foundation Trust

Contents

Summary of Renal Association Proposals	4
Introduction.....	5
Hypertension management for individuals without CKD.....	5
Measurement of BP – translating the evidence into clinical practice.....	6
Potential adverse effects associated with intensive BP lowering	7
Adherence testing	8
Dietary modifications	8
BP targets for individuals with pre-existing CKD	9
Implementing these proposals in a post-COVID-19 world	11
References	11

Method used to arrive at a recommendation

The recommendations for the first draft of this guideline resulted from a collective decision reached by informal discussion by the authors and, whenever necessary, with input from the Chair of the Clinical Practice Guidelines Committee. If no agreement had been reached on the appropriate grading of a recommendation, a vote would have been held and the majority opinion carried. However this was not necessary for this guideline.

Conflicts of Interest Statement

All authors made declarations of interest in line with the policy in the Renal Association Clinical Practice Guidelines Development Manual. Further details can be obtained on request from the Renal Association.

Acknowledgements

The committee would like to thank Mr Bruno Mafri and Ms Tina Dilloway for their valuable contribution to the section regarding lifestyle management.

Summary of Renal Association Proposals

Measuring blood pressure:

- 1.1 We propose that office-based blood pressure measurements using a standardized technique (e.g. <https://bihsoc.org/resources/bp-measurement/measure-blood-pressure>) should guide all treatment decisions.
- 1.2 We propose that non-office blood pressure measurements, taken using a standardised technique and validated monitor (see <https://bihsoc.org/bp-monitors/>) be considered a pragmatic alternative to office-based measurements.

Blood pressure targets for individuals without Chronic Kidney Disease:

- 2.1 We propose adopting a shared-decision making approach when considering blood pressure treatment and targets for any patient whenever possible.
- 2.2 We propose aiming for a blood pressure target of <130/80 mmHg in patients at elevated cardiovascular risk if, following a shared-decision making discussion, it is tolerated by the individual.
- 2.3 We propose that there is not enough evidence to inform on a blood pressure target in patients without cardiovascular risk factors, however if tolerated, aiming <130/80 mmHg is a reasonable goal.
2. We propose that for young adults, lowering blood pressure to the median value for age be considered.

Adherence testing:

- 3.1 We propose objective testing of medication adherence in those with resistant hypertension.

Dietary modifications:

- 4.1 We propose reducing dietary salt intake to a maximum of 6g/day and 3g/day by 2025.
- 4.2 We propose not using salt substitutes in place of dietary salt reduction.

Blood pressure target for individuals with Chronic Kidney Disease:

- 5.1 We propose aiming for a blood pressure target of <130/80 mmHg for all patients with chronic kidney disease, (irrespective of albuminuria), to improve cardiovascular outcomes if, following a shared-decision making discussion, it is tolerated by the individual.

Introduction

The publication in 2019 of NICE guideline NG136 ‘Hypertension in adults: diagnosis and management’¹ followed the updated guidance for the management of hypertension from the American College of Cardiology (published in 2017)² and the European Society of Cardiology/European Society of Hypertension (published in 2018).³ From a ‘nephrological’ perspective the American and European guidelines both make recommendations specific to the management of hypertension in patients with chronic kidney disease (CKD), whereas this issue was beyond the remit of the 2019 NICE guideline and instead readers are referred back to the (broader) 2014 NICE guideline⁴ for the general management of CKD.

Since 2014 a number of important studies have been published pertaining to the management of hypertension, both in general and in the context of CKD. The ‘SPRINT’ trial⁵ has received particular attention and has significantly influenced the guidance issued by both the Americans and the Europeans but impacted much less on the NICE recommendations.

Currently there is no UK guideline which has incorporated this newer evidence to assist those managing patients with CKD. Thus UK clinicians may be left with the dilemma of which guideline to follow. This document therefore has two objectives: first to provide a commentary on NICE guideline NG136 highlighting issues relevant to the renal community, including points of controversy and where the opinions of the Renal Association and NICE differ; secondly we provide (opinion-based) proposals for clinical practice regarding blood pressure (BP) management both in general and also specifically for patients with CKD considering the most recent evidence. We also consider the recent impact of COVID-19.

Hypertension management for individuals without CKD

The Renal Association supports the majority of recommendations in NICE guideline NG136 and considers the document pragmatic and clinically useful. Of the updated recommendations it is perhaps number 1.4.20 that has caused the most controversy:

1.4.20 Reduce clinic BP to below 140/90 mmHg and maintain that level in adults with hypertension aged under 80. [NICE, 2019]

This recommendation is in contrast to that in both the American and European guidelines which generally advocate (if tolerated) a BP target of under 130/80 mmHg. The guidance issued by these two bodies has been heavily influenced by the results from the SPRINT trial.⁵ It is interesting that SPRINT was also considered in depth by NICE but interpreted differently – ultimately with the conclusion that UK BP targets should not be changed from those recommended in the previous NICE guideline of 2016.

Briefly, SPRINT was a randomised trial of over 9000 patients (without diabetes but with other cardiovascular risk factors) with a baseline SBP of over 130 mmHg. Patients were assigned to intensive BP lowering (target SBP <120 mmHg) or standard treatment (target SBP <140 mmHg). BP was measured using a strict, standardised, office-based protocol (discussed in depth below). The mean achieved BP in the two groups

was 121 and 136mmHg respectively. The composite primary end-point of death from cardiovascular cause or a major cardiovascular event (myocardial infarction, acute coronary syndrome, stroke, or heart failure) was observed in 1.65% per year in the intensive arm vs. 2.19% per year with standard care – a hazard ratio of 0.75 in favour of intensive treatment and the trial was terminated early (after 3.3 years instead of the planned 5 years) due to the magnitude of effect.

So why, despite these strongly positive results, did NICE elect not to lower its recommended BP target? In the rationale and impact section of NG136, two main concerns are raised. The NICE committee question the applicability of the technique used to measure BP in SPRINT to everyday UK practice. In addition, the potential adverse effects from intensive BP lowering - in particular adverse renal events – seen in the trial are emphasised. The Renal Association’s viewpoint on these two issues differs somewhat from those held by NICE.

Measurement of BP – translating the evidence into clinical practice

In SPRINT, BP was measured using an automated device following five minutes of quiet rest and taking the mean of three readings. The SPRINT protocol did not specify whether BP readings were taken with healthcare workers ‘in attendance’ or not. The influence of attended vs. unattended BP measurements in SPRINT has been assessed.⁷ Interestingly (and perhaps surprisingly), the presence of a healthcare worker during BP measurement did not impact on the reading obtained. In its guideline, NICE has raised concern as to whether this ‘standardised’ BP measurement technique can be translated into the typical, busy outpatient clinic environment or GP practice. This issue was seemingly a strong influential factor in the decision made by NICE to not lower the recommended BP target despite the SPRINT results. This concern is despite NICE recommendation 1.1.4:

1.1.4 When measuring blood pressure in the clinic or in the home, standardise the environment and provide a relaxed, temperate setting, with the person quiet and seated, and their arm outstretched and supported. Use an appropriate cuff size for the person’s arm [NICE 2011, amended 2019].

All recent informative trials have used standardised BP measurement. The evidence-based approach, therefore, would be to issue clear guidance that casual office BP measurements should never be used to guide therapy, and that standardised office BP measurements (such as those detailed by the British and Irish Hypertension Society - <https://bihsoc.org/resources/bp-measurement/measure-blood-pressure>) should be offered to all patients in whom BP measurement is undertaken in a clinic setting. Other options, including home and 24 hour ambulatory BP monitoring (ABPM) are obviously available and whilst these have not been widely used in the trial setting to assess BP targets, pragmatically the Renal Association considers them to be reasonable alternatives and likely to provide useful information to guide therapy.

Home monitoring is likely to take on an even more important role in the future given the recently imposed constraints conferred by the COVID-19 pandemic. Therefore, a randomised trial of BP management utilising out-of-office measurements is urgently needed. NICE guideline NG136 does support the use of home monitoring and suggests a target of <135/85 mmHg (compared with the office target of <140/90 mmHg) although the evidence cited to support this recommendation is weak (essentially the mean difference

between office and home BPs at population level). The modality of BP monitoring will differ depending on patient choice and circumstances. However, whichever technique is employed, it needs to conform to the standard adopted in the trial setting in order for evidence-based targets to be relevant.

Renal Association proposals – measuring blood pressure

1.1 We propose that office-based blood pressure measurements using a standardized technique (e.g. <https://bihsoc.org/resources/bp-measurement/measure-blood-pressure>) should guide all treatment decisions.

1.2 We propose that non-office blood pressure measurements, taken using a standardised technique and validated monitor (see <https://bihsoc.org/bp-monitors/>) be considered a pragmatic alternative to office-based measurements.

Potential adverse effects associated with intensive BP lowering

It is perhaps unsurprising that in SPRINT, intensive BP lowering was associated with a greater risk of syncope, hypokalaemia and acute kidney injury (AKI). The risk of incident CKD (defined as a decline in eGFR >30% to an eGFR <60ml/min) was also increased with intensive BP control (1.21% per year versus 0.35% per year). However, no increase was seen in bradycardia or injurious falls. Interestingly, orthostatic hypotension was seen less frequently in the intensive arm.

The issue of AKI with more intensive BP control has been looked at in a separate sub-analysis of SPRINT participants.⁸ Whilst AKI was more common with intensive BP control, events were usually mild and reversible - likely reflecting in most cases the haemodynamic consequences of the commonly employed anti-hypertensive agents. Work demonstrating no increase in biomarkers of tubular damage from those in the more intensive BP lowering arm, despite lower eGFR, gives weight to this theory and is reassuring.⁹

There is very limited evidence relating to how patients may balance the risks versus benefits of aiming for a lower BP target. Shared-decision making is the key to this issue and individualising treatment goals and supplying patients with sufficient information to aid discussions, the optimal strategy. goals and supplying patients with sufficient information to aid discussions, the optimal strategy.

Renal Association proposals – blood pressure targets for individuals without CKD

2.1 We propose adopting a shared-decision making approach when considering blood pressure treatment and targets for any patient whenever possible.

2.2 We propose aiming for a blood pressure target of <130/80 mmHg in patients at elevated cardiovascular risk if, following a shared-decision making discussion, it is tolerated by the individual.

2.3 We propose that there is not enough evidence to inform on a blood pressure target in patients without cardiovascular risk factors, however if tolerated, aiming <130/80 mmHg is a reasonable goal.

2.4 We propose that for young adults, lowering blood pressure to the median value for age be considered.

Adherence testing

The Renal Association fully supports the emphasis NG136 places on assessment of medicines optimisation and adherence at all stages of hypertension treatment and in particular the reassessment of adherence in those reaching 'step 4' of the treatment pathway:

1.4.38 Before considering next step treatment for hypertension discuss with the person if they are taking their medicine as prescribed and support adherence in line with NICE's guideline on medicines adherence. [NICE, 2019]

1.4.45 Before considering further treatment for a person with resistant hypertension:

- **Discuss adherence (see recommendation 1.4.38). [NICE, 2019]**

The NICE guideline references the Medicines Adherence Guideline (CG76), however CG76 lists only generic methods for assessment of adherence such as patient self-reporting, using records of prescription re-ordering, pharmacy patient medication records, and return of unused medicines. Given the significant issue of medication non-adherence in patients with hypertension¹⁰ the Renal Association feel it would have been useful for NICE to discuss more objective strategies of direct adherence testing. These include observed pill taking with day case BP/ABPM monitoring, as well as urinary drug metabolite analysis which is now more widely available in the UK (Birmingham Heartlands and Leicester centres are supporting large number of hypertension centres across the country).

Renal Association proposal – adherence testing

3.1 We propose objective testing of medication adherence in those with resistant hypertension.

Dietary modifications

The lifestyle recommendations in NG136 remain essentially unchanged from the previous NICE guideline. Specifically, in terms of salt intake NICE recommendation 1.4.5 states:

1.4.5 Encourage people to keep their dietary sodium intake low, either by reducing or substituting sodium salt, as this can reduce blood pressure [NICE, 2004, amended 2019]

A footnote urges caution about using salt substitutes containing potassium chloride for certain people, including those with diabetes and/or kidney disease and those on blockers of the renin-angiotensin system.

Salt reduction in these groups is advised. The guideline may therefore be read as implying that substituting sodium salt is an alternative to reducing dietary sodium (salt).

A number of studies have showed an association between decreased salt intake and lower blood pressure^{11,12} and the Renal Association feel that the emphasis should be on reducing dietary salt intake for everyone (to the current target of 6g/day and 3g/day by 2025). The terminology ‘salt’ rather than ‘sodium’ would be preferable as this is recommended for food labelling and is more easily understood by the public.

Substituting sodium with potassium salts risks perpetuating a salty palate and it would be better to adapt to foods without salt and appreciate the natural flavours of food. The ‘DASH’ diet is not mentioned by NICE but we feel is worthy of note as it can help to lower blood pressure.¹³ Certain foods, in particular black liquorice, should be discouraged owing to its BP elevating potential.¹⁴ Providing primary health care professionals with training and/or support on how to provide dietary education would be valuable. Registered dietitians are best placed to provide this.

Renal Association proposals – dietary modifications

4.1 We propose reducing dietary salt intake to a maximum of 6g/day and 3g/day by 2025.

4.2 We propose not using salt substitutes in place of dietary salt reduction.

BP targets for individuals with pre-existing CKD

NICE guideline NG136 specifically *excludes* patients with CKD and readers are instead referred back to the 2014 NICE recommendations⁴ for general CKD management. This 2014 guideline offers two recommendations regarding BP control, namely:

1.6.1: In people with CKD aim to keep the systolic BP below 140 mmHg (target range 120–139 mmHg) and the diastolic BP below 90 mmHg [NICE, 2014].

1.6.2: In people with CKD and diabetes, and also in people with an ACR of 70 mg/mmol or more, aim to keep the systolic BP below 130 mmHg (target range 120–129 mmHg) and the diastolic BP below 80 mmHg [NICE, 2014].

The 2017 American Guidelines² recommend a lower BP target for all adults with CKD:

9.3: Adults with hypertension and CKD should be treated to a BP goal of less than 130/80 mmHg.

The 2018 European Hypertension guidelines³ offer a third alternative to the NICE and American recommendations:

8.12: In patients with diabetic or non-diabetic CKD it is recommended to lower SBP to a range of 130–139 mmHg. Individualized treatment should be considered according to its tolerability and impact on renal function and electrolytes.

The American and European guidelines for CKD have been influenced (to different degrees) by SPRINT and it is anticipated that the eagerly awaited KDIGO hypertension guidelines (presented in part at Kidney Week

2019) will follow suit. SPRINT included over 2500 patients with (non-diabetic) CKD with a mean eGFR of 48ml/min and minimal proteinuria (mean ACR 9mg/mmol). Of note, SPRINT excluded patients with proteinuria >1g/day, diabetes, polycystic kidney disease, and patients with eGFR <20ml/min.

Analysis of the main trial indicated that the presence of CKD did not significantly modify the primary outcome (p value for interaction = 0.36). In addition, a pre-specified subgroup analysis was undertaken for this CKD cohort.¹⁵ Interpretation of the data needs relative caution as the study was not powered for subgroup analysis and was terminated early, however the results are worthy of discussion. The primary composite cardiovascular outcome occurred in 112 of those in the intensive group and 131 in the standard group with a hazard ratio of 0.81 (95% confidence interval 0.63 to 1.05). A lower rate of all-cause death was seen in the intensive arm with a hazard ratio of 0.72 (95% confidence interval 0.53 to 0.99). The proportional reduction in CV events and total mortality was similar in the CKD subgroup and in the main trial (p for interaction 0.30 and 0.95 respectively).

In terms of renal disease progression, the pre-specified composite endpoint of >50% reduction in eGFR or occurrence of ESRD did not differ between groups although the number of events was extremely small (15 vs 16 in the intensive and standard arms respectively). A reduction in eGFR of >30% did occur more frequently with intensive BP lowering (as it did in the main trial) but whether this is clinically relevant given the reduction in 'harder' endpoints remains uncertain. Further analysis has suggested that the magnitude of effect in reducing cardiovascular endpoints with intensive BP reduction may be attenuated at lower levels of eGFR and this issue requires further study.¹⁶

The adverse effects observed in the CKD cohort essentially mirrored those in the main trial with increased risks of potassium disturbances and AKI in the intensive arm. However, no increase in hypotension-related events was seen including syncope and injurious falls. No differences were seen in sodium abnormalities or bradycardia.

Previous studies assessing optimal BP targets in CKD have primarily focussed on renal outcomes.¹⁷ These have differed by design and inclusion criteria and yielded varying results. None have been sufficiently powered to evaluate hard cardiovascular endpoints or mortality. Thus data from SPRINT provide the strongest evidence to date to inform about the effects of intensive BP lowering on cardiovascular events in patients with renal impairment. This is important considering the extremely high burden of cardiovascular-related morbidity and mortality in individuals with CKD.

Intensive lowering of BP in patients with CKD will undoubtedly lead to adverse sequelae in a minority of patients. Therefore, the importance of adopting a shared-decision making process regarding individualised BP targets cannot be emphasised enough. Monitoring BP using a standardised technique is also vital. Nonetheless the Renal Association considers that on balance, aiming for a BP of <130/80 mmHg is likely to be beneficial in the majority of individuals with CKD and treating to this target should be considered in all patients.

Renal Association proposal – blood pressure target for individuals with Chronic Kidney Disease

- 5.1 We propose aiming for a blood pressure target of <130/80 mmHg for patients with chronic kidney disease to improve cardiovascular outcomes if, following a shared-decision making discussion, it is tolerated by the individual.**

Implementing these proposals in a post-COVID-19 world

COVID-19 has thrown us into a clinical environment where the majority of outpatient services had to be reconfigured overnight into virtual clinics. This model of working was new for many clinicians, and has posed many new challenges in delivering safe patient care. There is a risk of clinician inertia in treating to lower BP targets that might be heightened by the challenges of virtual clinics being delivered without pre-designed pathways for service delivery. Obtaining reliable and accurate blood pressure readings, monitoring of eGFR and electrolytes following medication changes, and the inability to clinically examine patients in a remote virtual clinic are some of the challenges that we face when addressing hypertension management.

Clinical guidelines provide blood pressure targets based on clinic or office blood pressure readings. With a move towards virtual outpatient services there is an increasing need for research recommendations based on home blood pressure readings. It is crucial that home readings are obtained with validated and calibrated machines, in a standardised way, to ensure reliable and clinically meaningful interpretation. This requires supporting patients in choosing validated monitors (<https://bihsoc.org/bp-monitors/for-home-use/>), providing education about correct monitoring techniques through either online resources (<https://bihsoc.org/resources/bp-measurement/hbpm/>) or delivered during the consultation (e.g. supervised readings in first video consultation), and ability to record that data into the patient record (e.g. patient diaries emailed to clinical services, uploaded via Patient View, integrated into EPR / primary care record). There is an opportunity to improve engagement of patients with their hypertension management through home blood pressure readings, but clinicians also need to be mindful of the risk of suboptimal care being delivered for those without the resources (financial, language barriers, technological or telecommunication challenges etc.) to engage with virtual clinics.

References

1. National Institute for Health and Care Excellence: Hypertension in adults diagnosis and management (NG136) [Internet] 2019 August 28th; 1-40
2. 2017 *Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults*. J Am Coll Cardiol. Sep 2017, **23976**; DOI: 10.1016/j.jacc.2017.07.745
3. Williams B et al. 2018 *ESC/ESH Guidelines for the management of arterial hypertension*. European Heart Journal (2018) **39**, 3021–3104
4. National Institute for Health and Care Excellence: Chronic Kidney Disease in adults assessment and management (CG182) [Internet] 2014 July 23rd; 1-60
5. Wright JT Jr., Williamson JD, Whelton PK, et al. *SPRINT Research Group: A randomized trial of intensive versus standard blood pressure control*. N Engl J Med **373**: 2103–2116, 2015

6. National Institute for Health and Care Excellence: Hypertension in adults diagnosis and management (CG127) [Internet] 2011 August 24th
7. Johnson, KC, Whelton, PK, Cushman, WC et al. *Blood Pressure Measurement in SPRINT (Systolic Blood Pressure Intervention Trial)*. *Hypertension* 2018; **71**:848-857
8. Rocco, MV, Sink, KM, Lovato, LC et al. *Effects of Intensive Blood Pressure Treatment on Acute Kidney Injury Events in the Systolic Blood Pressure Intervention Trial (SPRINT)* *Am J Kidney Dis.* 2018 March ; **71**(3): 352–361.
9. Malhotra R, Craven T, Ambrosius WT, et al. *SPRINT Research Group: Effects of Intensive Blood Pressure Lowering on Kidney Tubule Injury in CKD: A Longitudinal Subgroup Analysis in SPRINT.* *Am J Kidney Dis* pii: S0272-6386 (**18**)30879-5, 2018,
10. Hameed MA, Dasgupta I. *Medication adherence and treatment resistant hypertension: a review.* *Drugs in Context* 2019; **8**: 212560
11. He FJ, Li J, Macgregor GA. *Effect of longer-term modest salt reduction on blood pressure.* *Cochrane Database Syst Rev.* 2013 Apr 30;**4**:CD004937.
12. Aburto NJ, Ziolkovska A, Hooper L, et al. *Effect of lower sodium intake on health: systematic review and meta-analyses.* *BMJ.* 2013 Apr 3;**346**
13. Bray GA, Vollner WM, Sacks FM, et al; *DASH Collaborative Research Group.* *A further subgroup analysis of the effects of the DASH diet and three dietary sodium levels on blood pressure: results of the DASH-Sodium Trial.* *Am J Cardiol.* 2004 Jul 15;**94**(2):222-7.
14. Sigurjonsdottir HA, Franzson L, Manhem K et al. *Liquorice-induced rise in blood pressure: a linear dose-response relationship.* *J Hum Hypertens* 2001; **15**:549–52. 10.1038/
15. Cheung AK, Rahman M, Reboussin DM, et al; *SPRINT Research Group: Effects of intensive BP control in CKD.* 2017. *J Am Soc Nephrol* **28**: 2812–2823
16. Obi Y, Kalantar-Zadeh K, Shintani A, et al. *Estimated glomerular filtration rate and the risk-benefit profile of intensive blood pressure control amongst nondiabetic patients: A post hoc analysis of a randomized clinical trial.* 2018 *J Intern Med* **283**: 314–327.
17. Chang AR, Loser M, Malhotra R et al. *Blood pressure goals in patients with CKD. A review of evidence and guidelines.* *Clin J Am Soc Nephrol* 2019; **14**: 1