

# Measures to Keep Staff Safe from COVID-19 in Haemodialysis Units



Staff are at risk from patients, staff and home contacts

Strong, supportive leadership



Clear communication




## SYMPTOMS-STAY AT HOME

- Symptomatic staff stay at home
- PCR test same day



## PPE – KEEP TRAINING UP TO DATE

- Mask, visor, gloves, gown/apron
- Dispose of, or clean, your visor
- Don't touch your mask



## SOCIAL DISTANCING

- In clinical areas and staff rooms
- Between computers/desks
- In meetings and handovers
- Stagger break times



## HAND HYGIENE

- In clinical areas
- In staff rooms
- In meeting rooms
- Audit compliance



## MAXIMISE FRESH AIR

- Understand ventilation system
- Increase intake of fresh air
- Keep windows open



## DECLUTTER AND CLEAN

- Appoint a tidy champion
- Clean workstations at least 4 times daily (keyboards, surfaces, computer mice, phones, etc.)



## MEETINGS & HANDOVERS

- Avoid face-to-face meetings
- Virtual meetings if possible
- Maintain social distancing
- Define max. no. people/room



## COVID-19 TESTING

- Patients** – PCR test weekly where community infection is high
- Staff** – regular lateral flow or PCR
- Measure staff sickness



## COVID-SAFE STAFF ROOMS

- Use own cup, don't share food
- Hand gel on entry and exit
- Keep mask on
- Increase cleaning



## VACCINATION

- Ensure all offered vaccine
- Discuss vaccine hesitancy
- Answer questions/concerns
- Re-offer vaccine



## INDIVIDUAL RISK & MENTAL HEALTH

- Identify those struggling or at risk
- It's OK not to be OK - discuss concerns, signpost to support
- Agree solutions



## RECOGNISE COVID OUTBREAKS

- 2 or more linked people in 14 days
- Test all staff and patients
- Report to local IPC team
- Perform root cause analysis