



Clinical practice guidelines for management of hypertension and renin- angiotensin-aldosterone system blockade in adults with diabetic kidney disease **2021 update**

Endorsed by:

Lead authors

Debasish Banerjee MD FRCP
Tahseen A Chowdhury MD FRCP
Indranil Dasgupta DM FRCP
Parijat De MD FRCP
Rosa Montero MD MRCP
Mona Wahba MA FRCP
Peter Winocour MD FRCP

ABCD-RA writing group additional members

Stephen Bain MD FRCP
Damian Fogarty MD FRCP
Andrew Frankel MD FRCP
Jannaka Karraliede MD FRCP
Patrick Mark MD FRCP
Ana Pokrajac MD FRCP
Adnan Sharif MD FRCP
Sagen Zac-Varghese MD MRCP

Authors

Stephen Bain, Professor of Medicine (Diabetes), Swansea University, Swansea
Debasish Banerjee, Consultant Nephrologist, St George's Hospital, London
Tahseen A Chowdhury, Consultant Diabetologist, Royal London Hospital, London
Indranil Dasgupta, Consultant Nephrologist, University Hospitals Birmingham NHS Foundation Trust
Parijat De, Consultant Diabetologist, City Hospital, Birmingham
Damian Fogarty, Consultant Nephrologist, Belfast Health and Social Care Trust, Belfast
Andrew Frankel, Consultant Nephrologist, Imperial College Healthcare NHS Trust, London
Janaka Karalliedde, Consultant Diabetologist, Guy's and St Thomas' Hospital London
Patrick Mark, Professor of Renal Medicine, University of Glasgow
Rosa M Montero, Consultant Nephrologist, Royal Berkshire NHS Foundation Trust, Reading
Ana Pokrajac, Consultant Diabetologist, West Hertfordshire Hospitals
Adnan Sharif, Consultant Nephrologist, University Hospitals Birmingham NHS Foundation Trust
Mona Wahba, Consultant Nephrologist, St Helier Hospital, Carshalton
Peter Winocour, Consultant Diabetologist, East and North Herts Institute of Diabetes and Endocrinology (ENHIDE), East and North Herts NHS Trust
Sagen Zac-Varghese, Consultant Diabetologist, ENHIDE, East and North Herts NHS Trust

Correspondence

Peter Winocour
Consultant Physician and Clinical Director for Diabetes and Endocrine Services
ENHIDE, Queen Elizabeth II Hospital, Welwyn Garden City, Herts AL7 4HQ
Email: peter.winocour@nhs.net
Phone: +44 (0)1438 288324

Citation for this document

Association of British Clinical Diabetologists and The Renal Association. *Clinical practice guidelines for management of hypertension and renin-angiotensin-aldosterone system blockade in adults with diabetic kidney disease*. London: ABCD, 2021.

Review date: March 2023

Copyright

All rights reserved. No part of this publication may be reproduced, distributed or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the written permission of the copyright owner. Applications to reproduce any part of this publication should be addressed to the publisher.

Copyright © Association of British Clinical Diabetologists 2021

Contents

Introduction	5
Evidence grades for the recommendations	5
Search strategy.....	5
Note on revisions	6
Classification	7
Blood pressure targets	7
eGFR and ACR categories	8
1 Hypertension management and renin-angiotensin-aldosterone system blockade in people with type 1 diabetes.....	9
Recommendations	9
Audit standards	10
Areas that require further research.....	10
Introduction	10
Hypertension in people with type 1 diabetes.....	11
Modulation of the RAAS in people with type 1 diabetes.....	14
2 Hypertension management and renin-angiotensin-aldosterone system blockade in people with type 2 diabetes, nephropathy and/or early CKD stages 1–3	19
Recommendations	19
Audit standards	19
Areas that require further research.....	20
Introduction	20
The renin-angiotensin-aldosterone system	21
Hypertension in people with type 2 diabetes.....	21
The role of home and ambulatory blood pressure measurement	22
Lifestyle modification and impact on blood pressure.....	22
Blood pressure lowering agents.....	22
Adherence with therapy	23
RAAS blockade in people with type 2 diabetes without proteinuria	23
RAAS blockade in people with type 2 diabetes and proteinuria or microalbuminuria	23
Use of dual blockade with ACEIs and ARBs in people with type 2 diabetic nephropathy ..	23
Aldosterone blockade in people with type 2 diabetic nephropathy	24
Use of direct renin inhibitors in diabetic nephropathy.....	26
When should RAAS blockade be stopped?	26
Other agents for blood pressure lowering in people with type 2 diabetes and nephropathy	27
3 Hypertension management and renin-angiotensin-aldosterone system blockade in people with type 2 diabetes, nephropathy and/or later stage CKD stages 4 and 5 (non-dialysis)	30
Recommendations	30
Audit standards	31
Research recommendations	31
Introduction	31
Identification and monitoring of people with diabetes and CKD stages 4 and 5	32
Target blood pressure in people with diabetes and CKD stages 4 and 5, with or without significant albuminuria (UACR >3 mg/mmol [>26.55 mg/g]).....	32
Renin-angiotensin system blockade for blood pressure control in people with diabetes and CKD stages 4 and 5	33
Management of hyperkalaemia with renin-angiotensin system blockade in people with diabetes and CKD stages 4 and 5	34

Non-pharmacological management of hypertension in people with diabetes and CKD stages 4 and 5.....	35
4 Hypertension management and renin-angiotensin-aldosterone system blockade in people with diabetes and CKD stage 5 on dialysis (5D)	36
Recommendations	36
Audit standards	36
Areas that require further research.....	37
Introduction	37
Blood pressure measurement in people with diabetes who are on haemodialysis.....	38
Target blood pressure in people with diabetes who are on dialysis	39
Volume control in people with diabetes who are on dialysis.....	40
Salt restriction in people with diabetes who are on dialysis	41
Declarations of interest	43
References.....	44

Introduction

A significant percentage of people with diabetes develop chronic kidney disease (CKD), and diabetes is also a leading cause of end-stage kidney disease.¹ More than a quarter of people who are on dialysis in the UK have diabetes.² Diabetic kidney disease (DKD) is associated with high morbidity and mortality, predominantly related to cardiovascular complications, and the progression to kidney failure requiring renal replacement therapy. Hypertension is a modifiable risk factor for cardiovascular complications and progression of CKD.³

Angiotensin converting enzyme inhibitors (ACEi) and angiotensin receptor blockers are established treatments to slow the progression of DKD and reduce cardiovascular events. Novel agents such as sodium glucose cotransporter-2 (SGLT-2) inhibitors, non-steroidal selective mineralocorticoid receptor antagonists and endothelin A receptor antagonists have recently been demonstrated to improve clinical outcomes and lower blood pressure, and are likely to be used in the routine management of DKD in the future.⁴

The scope of this update includes lifestyle advice, blood pressure targets and antihypertensive therapies in different categories of patients with DKD. Accurate measurement of blood pressure is vitally important before starting and during monitoring of antihypertensive treatment. Separate recommendation for blood pressure measurement has not been made in this guidance. Readers may follow the [British and Irish Hypertension Society's guidance](#) on standardised, automated blood pressure measurement. The blood pressure thresholds and targets in this guideline refer to standardised office blood pressure readings unless specified otherwise.

This guidance is for a variety of clinicians who treat people with diabetic kidney disease, including primary care physicians and specialists in diabetes, cardiology and nephrology. It intends to harmonise practices of blood pressure monitoring, and pharmacological and non-pharmacological management of hypertension, which vary considerably in different settings.

Evidence grades for the recommendations

The following evidence grading has been used to determine the strength of the recommendations; the suggested audit standards; and the questions for areas that require future research.

- 1A – Strong recommendation: high-quality evidence
- 1B – Strong recommendation: moderate-quality evidence
- 1C – Strong recommendation: low-quality evidence
- 1D – Strong recommendation: very low-quality evidence
- 2A – Weak recommendation: high-quality evidence
- 2B – Weak recommendation: moderate-quality evidence
- 2C – Weak recommendation: low-quality evidence
- 2D – Weak recommendation: very low-quality evidence

Search strategy

The recommendations are based on a review of literature carried out initially between October 2013 and December 2016 and a further review carried out in January 2020 for the current update. We searched the Cochrane Library, PubMed/MEDLINE, Google Scholar and Embase, using the following key terms: type 1 diabetes, type 2 diabetes, hypertension, albuminuria, microalbuminuria, microvascular complications, nephropathy, chronic kidney disease, angiotensin-converting enzyme inhibitors, angiotensin receptor blockers and mineralocorticoid antagonists.

Note on revisions

This guidance updates *Hypertension management and renin-angiotensin-aldosterone system blockade in patients with diabetes, nephropathy and/or chronic kidney disease* published in July 2017. For ease of recognition, key revisions and additions have been highlighted in pale yellow like this.

Classification

This guidance uses the following descriptions to differentiate and classify kidney disease. Diabetic kidney disease (DKD) is used throughout as a suitable umbrella term to include both diabetic nephropathy (DN) and diabetes mellitus and chronic kidney disease (DM CKD).

DKD is a progressive condition strongly associated with adverse cardiovascular and renal outcomes, including acute kidney injury (AKI), heart disease and mortality. A small but significant percentage of people may progress to end-stage kidney disease (ESKD), requiring renal replacement therapy.

Diabetic nephropathy (hyperglycaemia-induced glomerular disease) comprises a pattern of gradually rising urinary albumin:creatinine ratio followed by a steady reduction in estimated glomerular filtration rate (eGFR). However, hypertension and renal atheromatous disease often coexist in people who have diabetes and can cause varying degrees of damage. CKD can also arise from causes other than diabetes and vascular disease, therefore diabetes and CKD may coexist with no causal link (although this is rare). For practical purposes, people with type 2 diabetes and CKD can be classified as having DKD when no other basis for CKD is identified.⁴

Blood pressure targets

Table 1 Blood pressure targets in people with diabetes through stages of kidney function impairment

	Stage of kidney function impairment				
	Normal kidney function, normo-albuminuria	Normal kidney function, micro-albuminuria	CKD stages 1–3	CKD stages 4–5 (non-dialysis)	CKD stage 5 (dialysis)
Type 1 diabetes in mmHg (evidence grade)	<140/80–90 (2D) <120/80 (2D) [^]	≤130/80 (1B) 120/80 (2D) [^]	≤130/80 (1B) 120/80 (2D) [^]	≤140/90 (1B) ≤130/80 for albuminuric(2C)	≤140/90 (2D) ^{***} (interdialytic BP)
Type 2 diabetes in mmHg (evidence grade)	<140/90 (1D) <150/90 (2B) ^{**} (for ≥75 years)	<130/80 (2D)	<130/80 (2D)	<140/90 (1B) [*] <130/80 for albuminuric (2C)	<140/90 (2D) ^{***} (interdialytic BP)

CKD=chronic kidney disease; BP=blood pressure

*For adults >65 years a higher target >140/90 may be appropriate

**For frail adults >75 years a higher target >150/90 may be appropriate to avoid side effects

***Monitor and target interdialytic home BP for people on haemodialysis

[^] Lower targets for younger adults aged <30

eGFR and ACR categories

Fig 1 Renal Association classification of estimated glomerular filtration rates (eGFRs) and albumin:creatinine ratio (ACR) categories

GFR and ACR categories and risk of adverse outcomes			ACR categories (mg/mmol), description and range		
			<3 Normal to mildly increased	3–30 Moderately increased	>30 Severely increased
			A1	A2	A3
GFR categories (ml/min/1.73m ²), description and range	≥90 Normal and high	G1	No CKD in the absence of markers of kidney damage		
	60–89 Mild reduction related to normal range for a young adult	G2			
	45–59 Mild–moderate reduction	G3a ¹			
	30–44 Moderate–severe reduction	G3b			
	15–29 Severe reduction	G4			
	<15 Kidney failure	G5			

Increasing risk ↓

→ Increasing risk

¹ Consider using eGFR_{cystatinC} for people with CKD G3aA1 (see recommendations 1.1.14 and 1.1.15)

Abbreviations: ACR, albumin:creatinine ratio; CKD, chronic kidney disease; GFR, glomerular filtration rate

Adapted with permission from Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group (2013) KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. Kidney International (Suppl. 3): 1–150

1 Hypertension management and renin-angiotensin-aldosterone system blockade in people with type 1 diabetes

Recommendations

The following are recommendations for renin-angiotensin-aldosterone system (RAAS) blockade and hypertension management in people with type 1 diabetes.

- 1 In people with type 1 diabetes and urine albumin:creatinine ratio (UACR) <3 mg/mmol (<26.55 mg/g), we recommend a threshold for blood pressure therapy of a persistent upright (sitting or standing) blood pressure that is greater than or equal to 140/90 mmHg (Grade 1B).^{*}
In children and adolescents with type 1 diabetes, the threshold for high blood pressure is an average systolic blood pressure and/or diastolic blood pressure greater than the 95th percentile for the person's gender, age and height on more than three occasions (Grade 1B).[†]
- 2 We recommend that angiotensin-converting-enzyme inhibitor (ACEI) therapy should be used as a first-line agent for blood pressure lowering and, if ACEI therapy is contraindicated or not tolerated, angiotensin receptor blockers (ARBs) should be considered (Grade 1B).
- 3 In most adults with type 1 diabetes and persistent UACR >3 mg/mmol (>26.55 mg/g), we recommend that ACEI therapy should be considered irrespective of blood pressure, and that the target upright blood pressure should ≤130/80 mmHg in younger adults (Grade 1B), but ≤140/90 mmHg for those aged over 65 (Grade 2D). We recommend that the dose of ACEI should be titrated to the maximum tolerated (Grade 1B).
- 4 There is no current evidence to support a role for ACEI therapy for blood pressure control or renal protection in people with type 1 diabetes who are normotensive and have UACR <3 mg/mmol (<26.55 mg/g) (Grade 1C).
- 5 There is some evidence to support the use of candesartan to prevent the development or progression of retinopathy in people with type 1 diabetes who are normotensive and have UACR <3 mg/mmol (<26.55 mg/g) (Grade 1C).
- 6 There is no firm evidence to support a role of dual blockade of the RAAS in people with type 1 diabetes (Grade 1C).
- 7 We recommend that people with type 1 diabetes should be advised to stop RAAS-blocking drugs during periods of acute illness (Grade 1C).
- 8 We recommend that women of childbearing age should be encouraged to stop RAAS-blocking drugs prior to actively considering pregnancy (Grade 1B).

^{*} We suggest a target upright blood pressure in younger adults of 120/80 mmHg and 140/90 mmHg for those aged over 65 (Grade 2D).

[†] Between the ages of 30–65 years, for some people with higher lifetime risk through earlier age of onset of type 1 diabetes, it may be appropriate to target a diastolic BP of less than 80 mmHg (Grade 2C).

Audit standards

The following are suggested as audit standards for RAAS blockade and hypertension management in people with type 1 diabetes.

- 1 The proportion of people with type 1 diabetes, with micro- or macroalbuminuria who are treated with an ACEI or ARB at the maximum tolerated doses.
- 2 The proportion of people with type 1 diabetes with micro- or macroalbuminuria who are achieving a blood pressure of <130/80 mmHg for younger adults or <140/90 mmHg for those aged over 75.
- 3 The proportion of younger adults (under 30 years old) with persistent albuminuria who are achieving systolic blood pressure of <120 mmHg
- 4 The proportion of people with type 1 diabetes and albuminuria who are on submaximal doses of ACEIs or ARBs due to hyperkalaemia or adverse reactions.

Areas that require further research

The following areas lack good-quality evidence for RAAS blockade and hypertension management in people with type 1 diabetes, and hence further research is necessary.

- 1 In light of the fact that the presence of microalbuminuria in people with type 1 diabetes may not be the best predictor of whether they will develop progressive renal disease, what is the role for other markers (such as kidney injury molecule-1 (KIM-1)) in predicting the risk of renal disease in those with type 1 diabetes?
- 2 What is the role of dual RAAS blockade in people with type 1 diabetes and nephropathy?
- 3 What is the role of aldosterone receptor blockers or direct renin inhibitors in people with type 1 diabetes and nephropathy?
- 4 Is there a role for home or ambulatory blood pressure monitoring in the diagnosis and management of hypertension in people with type 1 diabetes, particularly in those who have diabetic autonomic neuropathy?
- 5 Does measurement of plasma renin activity have a role in screening and managing hypertension in people with type 1 diabetes?
- 6 Does tight glycaemic control and blood pressure lowering reduce the incidence of people developing microvascular complications?
- 7 What is the role of RAAS-blocking agents in people who have type 1 diabetes, progressive renal decline and normoalbuminuria?
- 8 What is the impact on renal function of lower blood pressure targets in younger people with type 1 diabetes and nephropathy?

Introduction

Despite improvements in prognosis, diabetic nephropathy in people with type 1 diabetes remains a major cause of end-stage kidney disease.⁵ The onset of micro- and macroalbuminuria in people with type 1 diabetes heralds not only an increased risk of renal disease, but also an increased risk of cardiovascular disease.⁶ Early prospective studies suggested that around 30–50% of people with type 1 diabetes will develop microalbuminuria, in whom a 6% increase in risk of coronary heart disease is seen per 5 mg increase in 24-hour albumin excretion rate (AER).⁶ The natural history of diabetic nephropathy in people with type 1 diabetes has, however, changed over the past 4 decades.

Studies in the 1970s and 1980s suggested that progression to end-stage kidney disease in people with macroalbuminuria would occur within 7 years.⁷ More recent follow-up data of significant numbers of people with type 1 diabetes suggest that end-stage kidney disease occurs in around 3% of people who have had diabetes for 10 years⁸ and in around 8% of people who have had diabetes for 30 years.⁹

There is a significant body of evidence to suggest that over-activation of the RAAS plays a major role in the pathogenesis of diabetic nephropathy in people with type 1 diabetes.¹⁰ Over-activation of the RAAS is observed in people with type 1 diabetes, even in the absence of diabetic nephropathy.¹¹ Angiotensin II-mediated increase in intraglomerular pressure appears to be an important mechanism by which renal disease progresses in people with type 1 diabetes who have diabetic nephropathy,^{12,13} and reductions in intraglomerular pressure may ameliorate glomerular injury. Angiotensin II also has mitogenic effects that may lead to mesangial expansion that is characteristic of diabetic nephropathy.¹⁴ Over-activation of the RAAS may be mediated by hyperglycaemia,¹⁵ and blockade of the RAAS may in turn be impaired by hyperglycaemia.¹³ RAAS over-activation is also described in people with type 1 diabetes who have glomerular hyperfiltration.¹⁶

Hypertension in people with type 1 diabetes

Risk factors for the development of nephropathy in people with type 1 diabetes include increasing age, duration of diabetes, male gender and hyperglycaemia.¹⁷ The possible role of genetic factors has long been hypothesised, due to the observation that a family history of hypertension appears to predict the development of nephropathy.¹⁸ It has been suggested that a family history of hypertension could be the basis for more intensive antihypertensive therapy in people with type 1 diabetes.

The risk factor that has the strongest association with progression of diabetic nephropathy is hypertension. Prospective evaluation of 148 people with type 1 diabetes who were normoalbuminuric showed that those who developed microalbuminuria had a significantly higher baseline blood pressure compared with those who remained normoalbuminuric (138/82 mmHg versus 123/73 mmHg).¹⁹ Similarly, follow-up of a Scandinavian cohort of over 300 children and adolescents with type 1 diabetes showed that systolic blood pressure was a major risk factor for the development of microalbuminuria over 5 years.²⁰ Analysis of 1,441 people with type 1 diabetes in the Diabetes Control and Complications Trial (DCCT) cohorts suggests that systolic blood pressure levels below 120 mmHg are associated with a 41% reduction in macroalbuminuria (95% confidence interval [CI] 5% to 63%) and a 68% reduction in CKD stage 3 (95% CI 25% to 84%).²¹ The Pittsburgh EDC study 25-year follow-up results support an optimal blood pressure of 120/80 in childhood onset type 1 diabetes.²²

The threshold for diagnosis of hypertension in people with type 1 diabetes varies according to national and international guidelines. The Kidney Disease Improving Global Outcomes (KDIGO) guidelines suggest a blood pressure goal of less than or equal to 140/90 mmHg if urinary AER is under 30 mg per 24 hours, or less than or equal to 130/80 mmHg if AER exceeds 30 mg per 24 hours, although they do not distinguish between type 1 and type 2 diabetes.²³ National Institute for Health and Care Excellence (NICE) guidelines on the management of people with type 1 diabetes suggest a blood pressure target of 130/80 mmHg in a patient with albuminuria (135/85 mmHg in those who were normoalbuminuric).²⁴ The American Diabetes Association and American Society of Nephrology consensus guidelines suggest that a blood pressure of less than 140/90 mmHg should be attained in all individuals with diabetes and renal disease, but they do not suggest a lower target and they do not distinguish between type 1 and type 2 diabetes.²⁵ In children with type 1 diabetes, the International Society for Pediatric and Adolescent Diabetes (ISPAD) defines hypertension as average systolic blood pressure and/or diastolic blood pressure that

is greater than the 95th percentile for the person's gender, age and height on more than three occasions, and suggests a target blood pressure of 130/80 mmHg.²⁶ Given the younger age of many adults with type 1 diabetes and the consequent longer lifetime blood pressure burden, we support the targets of 120–140/80 mmHg according to age and the presence or absence of albuminuria, with those aged over 65 being better suited to the 140/90 mmHg target.^{27,28,29}


The role of home and ambulatory blood pressure measurement in the diagnosis and management of hypertension in people with type 1 diabetes and nephropathy is unclear. Small cohort studies of children and adults with type 1 diabetes suggest that an increase in nocturnal systolic blood pressure or blunting of nocturnal dipping is an important factor in progression to microalbuminuria in people with type 1 diabetes.^{30,31} Due to a lack of robust evidence, no guidelines currently recommend ambulatory or home blood pressure monitoring to diagnose or manage hypertension in people with type 1 diabetes, although the ISPAD guidelines suggest that there may be a role for 24-hour blood pressure monitoring in the diagnosis of hypertension in children.²⁶

The importance of lifestyle measures (weight loss and salt intake reduction) are highlighted by a number of guidelines, and indeed a study suggests that lower sodium intake may improve the efficacy of RAAS blockade.³²

There is evidence to suggest that management of blood pressure in people with type 1 diabetes may be suboptimal. In a large cross-sectional study of people with type 1 diabetes in Scandinavia, patients on antihypertensive therapy who were achieving a blood pressure below 130/80 mmHg varied according to degree of albuminuria.³³ Blood pressure above 130/80 mmHg was seen in 74.6% of people who were normoalbuminuric; 71.2% of people who were microalbuminuric; 80.0% of people who were macroalbuminuric; 88.1% of people who were treated with dialysis; and 90.4% of people who had received a renal transplant.


An important point to consider is the presence of postural hypotension in people with type 1 diabetes. Autonomic neuropathy is often associated with postural hypotension, and people with type 1 diabetes should have their supine and standing blood pressure checked. A significant drop in blood pressure on standing (>20 mmHg) might alert the clinician to ensure that care is taken not to treat the person's blood pressure over-aggressively. Otherwise we advocate the use of upright (sitting or standing) blood pressure as the target blood pressure, as per [British and Irish Hypertension Society guidelines](#).³⁴

Figure 2: Steps in the management of hypertension in people with diabetes and CKD stages 4 and 5



Box 1 Lifestyle modification and impact on blood pressure

There is good evidence from a number of observational studies and randomised controlled trials that salt intake, weight and body mass index (BMI), exercise frequency and alcohol intake all have a significant impact on blood pressure levels.^{35–38}



Salt intake

The evidence base for the benefit of salt restriction in type 1 diabetes without advanced CKD is not strong. Reduced blood pressure has been found in some but not all short-term studies, but an important long-term observational study recorded higher dietary sodium intake was associated with higher all-cause mortality and the development of ESKD.^{39,40} The KDIGO guidelines suggest lowering salt intake to less than 90 mmol of sodium per day (<2 g of sodium, which corresponds to 5 g of sodium chloride).²³ High salt intake has a greater impact on blood pressure for people with diabetes, especially in those with CKD, due to their reduced ability to excrete salt load in their urine. Restricting salt intake lowers blood pressure by a moderate amount, as shown in a systemic review of seven trials where salt intake was restricted to 4–6 g (70–100 mmol), systolic blood pressure was reduced by 4.7 mmHg and diastolic blood pressure was reduced by 2.5 mmHg.⁴¹

Given that salt restriction is inexpensive and it helps to lower blood pressure in the general population, despite a lack of availability of large-scale, long-term randomised controlled trials of salt restriction in people with CKD, there is no reason to believe that it would not be beneficial, although it would add to the dietary restrictions for managing diabetes. A low-salt diet has been shown to reduce blood pressure and albuminuria in the short term in people who are on angiotensin receptor blockers (ARBs) and it may be a consideration for those with high blood pressure who have had a poor response to ACEIs or ARBs.^{42,43}

Weight and BMI

Although abdominal obesity has been associated with higher blood pressure and use of antihypertensive therapy in type 1 diabetes,⁴⁴ there is a dearth of evidence that weight reduction in type 1 diabetes reduces blood pressure, although this would be expected intuitively.⁴⁵ There is evidence of weight gain accompanied by increases in blood pressure in type 1 diabetes as a consequence of improved blood glucose control. The KDIGO guidelines recommend achieving or maintaining a healthy weight (BMI 20–25).²³ Some observational studies, but not randomised trials, suggest that weight loss is likely to improve blood pressure in people with CKD, but there is a lack of high-quality randomised controlled trials in this area.

Although obesity has been proposed to be a potential mediator of CKD progression, trials are conflicting and reliable data remain sparse. There is no role of weight loss diets in CKD either. Overall, achieving a healthy body weight will improve blood pressure levels and prognosis in CKD, particularly in the early stages (stages 1–2). Malnutrition needs to be avoided in more advanced stages of CKD.⁴⁶

Exercise programme

There is documentation that exercise training for 12 weeks or more reduces blood pressure in type 1 diabetes.⁴⁷ The KDIGO guidelines recommend undertaking an exercise programme that is compatible with cardiovascular health and tolerance, aiming for at least 30 minutes of exercise five times per week.²³ Increased physical exercise has a broad range of positive health outcomes in the general population. However, there are no randomised controlled trials in the CKD population; there are mostly observation studies. The benefits of exercise on blood pressure and on general health are likely to be similar in the CKD population as they are in the general population.⁴⁸

Alcohol intake

Evidence that alcohol intake affects blood pressure and reduction in intake helps blood pressure in type 1 diabetes is sparse. The KDIGO guidelines suggest limiting alcohol intake to no more than two standard drinks per day for men and no more than one standard drink per day for women.²³ Most of the effects of alcohol reduction are related to its effect on blood pressure; that is, suggesting that restricting alcohol intake would lower blood pressure. All the trial evidence is mostly related to the general population and there are no specific data on people with CKD, but the effects of alcohol intake on blood pressure are expected to be similar.⁴⁹

Modulation of the RAAS in people with type 1 diabetes

a Normoalbuminuria

There has been some interest in the use of agents that block the RAAS in the primary prevention of diabetic kidney disease. The use of ACEIs has been tested in people who are normotensive and normoalbuminuric, and there is little evidence of a protective effect on the development of diabetic nephropathy. Importantly, however, many of these studies have used definitions of blood pressure that would now be considered to be too high. A multicentre European study examined 530 people with type 1 diabetes and blood pressure under 155/90 mmHg.⁵⁰ The study found that during 2 years of treatment with lisinopril versus placebo, the ACEI showed no protective effect against the development of microalbuminuria. Similar findings have been shown with candesartan.⁵¹ Furthermore, a renal biopsy study of 285 people with type 1 diabetes who were normotensive and normoalbuminuric showed no effect of enalapril or losartan in the development of renal lesions.⁵² One short study did suggest a significant reduction in albumin:creatinine ratio (ACR) in 89 individuals with type 1 diabetes who were normotensive and normoalbuminuric and who were treated with placebo or perindopril for 4 months.⁵³ Currently however, the use of ACE inhibition in people with type 1 diabetes who are normotensive and normoalbuminuric cannot be recommended on the basis of trial evidence.

b Microalbuminuria

The onset of microalbuminuria in someone with type 1 diabetes was once thought to lead to inevitable progression to macroalbuminuria and thence to progressive kidney disease.⁵⁴ More recently, however, it has become clearer that microalbuminuria may remit in up to 40% of people with type 1 diabetes.^{55–57} In addition, microalbuminuria may not progress to macroalbuminuria in a significant number of people.⁵⁸ In the Renin-Angiotensin System Study, onset of microalbuminuria correlated poorly with renal biopsy findings of diabetic glomerulopathy.⁶⁰ Previous studies have also described progressive renal impairment without microalbuminuria in people with type 1 diabetes.^{59,60}

In adolescents with type 1 diabetes, modest but persistent elevations of UACR in the normal range may be associated with progression to persistent microalbuminuria.⁶¹ Furthermore, a persistently raised UACR at the upper limit of the reference range in adolescents is

associated with increasing aortic intima-media thickening, which is a sensitive marker of early atherosclerosis.⁶²

It is, however, recognised that the presence of microalbuminuria may not be the ideal risk marker for progressive renal dysfunction in people with type 1 diabetes.⁶³ Microalbuminuria may progress, stabilise or regress, and factors that govern this change are unclear, especially in adolescents and young adults who have improved glycaemia control.⁵⁶ Therefore, more reliable biomarkers or genetic markers are needed to predict which individuals are at the greatest risk of progressive renal disease. Many studies have looked at putative genetic loci within the RAAS for a genetic predisposition to diabetic nephropathy, but no clear correlation with nephropathy risk has been found in most studies.⁶⁴ Some authorities suggest that determination of serum cystatin C in people with diabetes and proteinuria may provide better risk stratification of subsequent end-stage kidney disease than determination of serum creatinine.⁶⁵ Serum concentration of tumour necrosis factor (TNF) receptors 1 or 2 (TNFR1, TNFR2) may also be predictors of future development of CKD stage 3 in people with type 1 diabetes.⁶⁶ If those who are at high risk of progression to diabetic nephropathy could be identified early, more intensive systematic therapy could be considered, for example pancreas transplantation.⁶⁷

There are few long-term studies that suggest that treating people with type 1 diabetes, microalbuminuria and normal blood pressure reduces end-stage kidney disease. There are, however, more short-term studies that focus on a change in AER rather than a change in renal function. A multicentre European study examined 79 individuals with microalbuminuria and blood pressure below 155/90 mmHg, and found a significant reduction in AER in the group of people who were treated with lisinopril compared with the patients who were treated with a placebo (-34.2 mg/min).³³ In an 8-year follow-up of a small number of people with type 1 diabetes and microalbuminuria, 10% of people who were treated with captopril progressed to macroalbuminuria, compared with 40% of those who were treated with a placebo;⁶⁸ therefore, treatment with captopril was associated with a reduction in progression of renal disease. Similarly, the Microalbuminuria Collaborative Study Group treated 235 people with microalbuminuria and blood pressure less than 160/90 mmHg with placebo or captopril.⁶⁹ Progression to macroalbuminuria was seen in 21.9% of the placebo-treated group compared with 7.2% of the captopril-treated group (the risk reduction was 69%). The Ace-Inhibitor Trial to Lower Albuminuria in Normotensive Insulin-Dependent Subjects (ATLANTIS) study used ramipril versus placebo in 140 patients with type 1 diabetes and microalbuminuria and normotension, and it showed regression to normoalbuminuria in 20% of the ramipril-treated group, compared with 4% of the placebo-treated group.⁷⁰ A further study of 20 individuals with type 1 diabetes, microalbuminuria and normal blood pressure who were treated with placebo or enalapril showed a reduction in progression to macroalbuminuria and a significant number of people regressed to normoalbuminuria.⁷¹

RAAS blockade may have positive longer-term impacts on renal haemodynamics in people with type 1 diabetes even when therapy is stopped. In the 5-year Renin-Angiotensin System Study, patients who were on RAAS blockade during the trial, but who stopped therapy after the trial, showed significantly greater renal haemodynamic responses to clamped hyperglycaemia and flow-mediated vasodilatation, which suggests that RAAS blockade has sustained, long-term protective effects.⁵²

In people who are hypertensive and have microalbuminuria, however, there is relatively strong evidence to suggest that ACE inhibition slows progression of diabetic nephropathy in people with type 1 diabetes and microalbuminuria.⁷² Meta-analysis suggests that ACEIs reduce progression of microalbuminuria to macroalbuminuria (odds ratio 0.38; 95% CI 0.25

to 0.57).⁷³ Outcomes in terms of the prevention of end-stage kidney disease, however, have not been reported.

c Macroalbuminuria

For decades, the presence of macroalbuminuria in people with type 1 diabetes has been considered to be a stage of irreversible kidney disease. However, recent long-term follow-up of 159 individuals with type 1 diabetes in the Diabetes Control and Complications Trial / Epidemiology of Diabetes Interventions and Complications (DCCT/EDIC) study showed that 10 years after onset of macroalbuminuria, the cumulative incidence of reduction to microalbuminuria was 52%.⁵⁸ The cumulative incidence of CKD stage 3 (estimated glomerular filtration rate (eGFR) <60 mL/min/1.73 m²) was 32%, and the cumulative incidence of end-stage kidney disease was 16% after 10 years, with better glucose and blood pressure control being the main factors associated with a lower risk of CKD progression. Therefore, while macroalbuminuria appears to be an important renal disease risk marker, it is far from inevitable that relentless progression to end-stage kidney disease will occur.

Seminal studies in the 1980s suggested that early aggressive antihypertensive therapy could reduce the rate of decline of renal function in people with diabetic nephropathy,⁷⁴ and further studies of people with type 1 diabetes, hypertension and macroalbuminuria demonstrated the specific protective effects of ACEI drugs on progression of albuminuria and renal disease.^{75–77} Meta-analysis of these studies suggests a long-term beneficial effect on preventing doubling of serum creatinine and development of end-stage kidney disease.⁷⁸

d Use of other agents that modulate the RAAS

Candesartan has been studied in people with type 1 diabetes and diabetic retinopathy.^{15,79} A beneficial effect of candesartan was seen in the progression of retinopathy; although in one study, this was limited to people with poor glucose control (glycated haemoglobin (haemoglobin A1c) >7.5%).¹⁵ Studies using angiotensin II receptor blockers (ARBs) have not been widely reported in type 1 diabetes and nephropathy. In a small Danish study, losartan was seen to attenuate AER in people with type 1 diabetes.⁸⁰ In the Renin-Angiotensin System Study, however,⁵² use of losartan did not appear to protect people from developing microalbuminuria: indeed 17% of patients on losartan developed microalbuminuria compared with 6% on a placebo and 4% on enalapril over 5 years.

It has been suggested that aldosterone escape during long-term RAAS blockade may be a mechanism by which ACE inhibition fails to prevent progressive renal disease in people with type 1 diabetes.⁸¹ Thus the use of aldosterone antagonists in such patients may be useful. Spironolactone has been investigated in a small study of people with type 1 diabetes and microalbuminuria.⁸² Spironolactone added to standard antihypertensive therapy reduced AER by 60%, with no drop in blood pressure and a minor drop in eGFR, although severe hyperkalaemia was seen in a small number of individuals.

Aliskiren, the direct renin inhibitor, has been trialled in people with type 2 diabetes and diabetic nephropathy, and no significant effect on renal outcomes has been noted, although a reduction in AER has been noted.⁸³ A small study of people with type 1 diabetes who were treated with aliskiren showed positive effects on renal haemodynamic indices and systemic vascular responses.⁸⁴ Furthermore, dual blockade with ACEI also showed beneficial effects on arterial compliance, flow-mediated dilatation and renal vasodilatation.⁸⁵ Further study of this group of drugs in type 1 diabetes is warranted.

Early studies of beta-blockade in people with diabetic nephropathy and type 1 diabetes suggest an equivalent effect to ACEI.⁸⁶ There is some suggestion that non-dihydropyridine

calcium channel blockade may have some of the benefits of dihydropyridine calcium channel blockers in the management of diabetic nephropathy.⁸⁷

Through their ability to reduce intraglomerular pressure, blood pressure and uric acid levels, sodium glucose cotransporter-2 (SGLT-2) inhibitors may offer the possibility of renal protection. One study suggests that SGLT-2 inhibitors can offer a reduction in glomerular hyperfiltration.⁸⁸ Recent analysis of the Empagliflozin, Cardiovascular Outcomes and Mortality in Type 2 Diabetes (EMPA-REG) study suggests significant renoprotection.⁸⁹ There is evidence from the Tandem 1 and 2 studies that sotagliflozin can reduce blood pressure and UACR over 12 months, with similar renal haemodynamic changes seen with SGLT-2 inhibition in type 2 diabetes.⁹⁰ However, the risk that these agents can cause ketoacidosis when they are given to people with type 1 diabetes may limit their potential use in this group.⁹¹

e Use of dual blockade in people with type 1 diabetes and diabetic nephropathy

Meta-analysis of a number of these studies of people with type 2 diabetes and nephropathy suggest a reduction in proteinuria, but at the expense of an increased risk of severe hyperkalaemia and episodes of acute kidney injury (AKI).^{92–94} More recently, however, a large randomised controlled trial involving people with type 2 diabetes suggests that RAAS dual blockade has no benefit in terms of mortality, but that it may increase the risk of hyperkalaemia and AKI.⁹⁵

It is currently unclear whether there is a role for dual blockade in people who have type 1 diabetes and a normal eGFR ($>60 \text{ mL/min/1.73 m}^2$) in whom albuminuria is uncontrolled or increasing. While this may reduce albuminuria, there is no evidence of a reduction in other renal or cardiovascular endpoints.

In type 1 diabetes the pathogenic processes that occur in the development and progression of diabetic nephropathy may be very different. Use of ACEIs is associated with a compensatory increase in plasma renin activity, and this effect may be ameliorated by the use of ARB drugs. There are, however, few studies on the use of dual RAAS blockade and outcomes in type 1 diabetes. One small study from India of 30 people who were treated for a short period with telmisartan and ramipril resulted in a reduction in ACR and blood pressure, with a slightly increased risk of hyperkalaemia.⁹⁶ A further small study of 21 people with type 1 diabetes showed that the addition of irbesartan to ACEI therapy in people with type 1 diabetes resulted in a 37% reduction in AER, along with significant reductions in blood pressure.⁹⁷ Further studies of dual RAAS blockade in type 1 diabetes are needed.

f When should RAAS blockade be stopped?

The use of RAAS-blocking drugs in early pregnancy has been associated with harm to the fetus, including cardiovascular, neurological and renal malformations,⁹⁸ although more recent surveys do not suggest that there is a high risk of these problems occurring.⁹⁹ Pregnancy is associated with a high risk of progression of diabetic nephropathy in people with type 1 diabetes, and the benefits of RAAS blockade in such individuals may outweigh the risks, but current advice is that RAAS-blocking drugs should be stopped when pregnancy is confirmed, and indeed when pregnancy is planned.

Drugs that block the RAAS reduce intraglomerular pressure and may lead to a rise in serum creatinine of up to 30%, which should then stabilise.¹⁰⁰ Some studies suggest that clinically significant renal artery stenosis may be quite common among people with diabetes, especially those with type 2 diabetes.¹⁰¹ While the use of drugs that modulate the RAAS may increase the risk of deterioration in renal function in people with renovascular disease, in practice such deterioration is rare.^{102,103}

RAAS blockade can lead to hyperkalaemia, which may be managed by dietary methods, diuretics or use of sodium bicarbonate. However, if the hyperkalaemia is severe and refractory to these measures, RAAS blockade may need to be stopped or reduced.¹⁰⁴ A further possible clinical scenario is someone with type 1 diabetes having deteriorating renal function despite having well-controlled blood pressure on ACEI drugs. Once renal dysfunction continues to escape, despite optimal therapy, a decision may need to be made about cessation of ACEI therapy, especially if there may be a degree of ischaemic nephropathy, renovascular disease or postural hypotension. RAAS blockade may also increase the risk of AKI in people with diabetes, and advice to stop these drugs during periods of acute illness should be considered.¹⁰⁵

2 Hypertension management and renin-angiotensin-aldosterone system blockade in people with type 2 diabetes, nephropathy and/or early CKD stages 1–3

Recommendations

The following are recommendations for renin-angiotensin-aldosterone system (RAAS) blockade and hypertension management in people with type 2 diabetes, nephropathy and/or early chronic kidney disease (CKD).

- 1 In people with type 2 diabetes and hypertension, we recommend salt intake of <90 mmol per day (<2 g per day of sodium – equivalent to 5 g of sodium chloride) (Grade 1C).
- 2 In people with type 2 diabetes, CKD and urine albumin:creatinine ratio (UACR) <3 mg/mmol (<26.55 mg/g), we recommend that their target upright blood pressure should be <140/90 mmHg, using antihypertensive therapy in the maximum tolerated doses (Grade 1D).
- 3 In people with type 2 diabetes, CKD and UACR >3 mg/mmol (>26.55 mg/g), we suggest aiming for a target upright blood pressure that is consistently <130/80 mmHg, using antihypertensive therapy in the maximum tolerated doses (Grade 2D).
- 4 There is no evidence to support either ACEI or ARB therapy as first-line blood pressure lowering agents in comparison with other antihypertensive agents in people with type 2 diabetes, normal renal function and normal UACR (<3 mg/mmol [<26.55 mg/g]) (Grade 1A).
- 5 We suggest that ACEIs (or ARBs if ACEIs are not tolerated) should be preferentially used in people with type 2 diabetes and CKD who have UACR >3 mg/mmol (>26.55 mg/g). We recommend that the dose of ACEI (or ARB) should be titrated to the maximum tolerated (Grade 2D).
- 6 There is currently no evidence to support the role of home or ambulatory blood pressure monitoring in people with type 2 diabetes and CKD stages 2 and 3 (Grade 1D).
- 7 There is currently no evidence to support the role of dual blockade of the RAAS in people with type 2 diabetes and CKD stages 1 to 3 (Grade 1B).
- 8 Upright blood pressure targets should be set at no lower than 150/90 mmHg in those with type 2 diabetes who are aged 75 years or over (Grade 2B).
- 9 We recommend that people with type 2 diabetes should be advised to stop RAAS-blocking drugs during periods of acute illness and restarted 24–48 hours after recovery from the illness (Grade 1C).

Audit standards

The following are suggested as audit standards for RAAS blockade and hypertension management in people with type 2 diabetes, early CKD and/or proteinuria.

- 1 The percentage of people with type 2 diabetes who have CKD and urine AER of >30 mg per 24 hours (UACR >3 mg/mmol [>26.55 mg/g]) who are achieving the target upright blood pressure of <130/80 mmHg.

- 2 The proportion of people with type 2 diabetes who have CKD and urine AER >30 mg per 24 hours (UACR >3 mg/mmol [>26.55 mg/g]) who are on ACEIs or ARBs.
- 3 The percentage of people with type 2 diabetes who have CKD and UACR >3 mg/mmol (>26.55 mg/g) who are not on ACEIs or ARBs (or who are on a submaximal dosage) due to hyperkalaemia or a decrease in estimated glomerular filtration rate (eGFR) of >25% or an increase in serum creatinine of >30%.
- 4 The number of people with type 2 diabetes and CKD who are on dual blockade of the RAAS.

Areas that require further research

The following areas lack good-quality evidence for RAAS blockade and hypertension management in people with type 2 diabetes, nephropathy and/or early CKD, and hence further research is necessary.

- 1 What is the best method for blood pressure measurement in people with type 2 diabetes who have CKD?
- 2 What is the evidence-based lower limit for blood pressure reduction in people with type 2 diabetes who have CKD?
- 3 Does reduction in albuminuria with agents that do not modify blood pressure improve hard cardiovascular and renal outcomes?
- 4 Should RAAS inhibition be maximised in people with CKD?
- 5 Can potassium binders enable a higher dosage of RAAS inhibitors with better attainment of blood pressure control and reduction in hyperkalaemia?
- 6 What are the best second- and third-line blood pressure lowering agents in people with type 2 diabetes who have CKD and proteinuria?
- 7 Is there a need for long-term outcome studies of non-dihydropyridine calcium channel blockers in diabetic nephropathy?

Introduction

Diabetic nephropathy is the leading cause of end-stage kidney disease and it is characterised by a triad of persistent albuminuria, hypertension and a decline in glomerular filtration rate (GFR). The presence of diabetic nephropathy increases cardiovascular morbidity and mortality and also increases progression to end-stage kidney disease.^{106–108} After approximately 20–25 years, 40% of people with type 2 diabetes develop evidence of diabetic nephropathy.¹⁰⁹ Both hypertension and hyperglycaemia are strong risk factors in determining progression of end-stage kidney disease and cardiovascular complications in diabetic nephropathy. Microalbuminuria is one of the earliest manifestations of kidney disease in people with diabetes and it predicts increased cardiovascular morbidity and mortality in people with both type 1 and type 2 diabetes.^{110,111} The prevalence of microalbuminuria in people who have had type 2 diabetes for 10 years is 25%, with an annual rate of progression to overt nephropathy of approximately 3%.¹⁰⁹

The risk of new as well as progressive microalbuminuria is significantly associated with high blood pressure.¹¹² In people with diabetes, cardiovascular and renal outcomes are adversely affected by the presence of hypertension and albuminuria.¹¹³ Thus controlling blood pressure and reducing albuminuria are important treatment goals in diabetic nephropathy. Baseline blood pressure levels have been shown to be a powerful determinant of subsequent kidney failure in large population-based studies.^{114,115} Unlike those with type 1

diabetes, a high proportion of people with type 2 diabetes often have microalbuminuria and overt nephropathy at diagnosis. Without intervention, 20–40% of people with type 2 diabetes and microalbuminuria will progress to overt nephropathy. After 20 years of overt nephropathy, approximately 20% of those people will progress to end-stage kidney disease.

The renin-angiotensin-aldosterone system

Dysregulation of the renin-angiotensin-aldosterone system (RAAS) plays a vital role in the pathogenesis of diabetic nephropathy, including pathogenesis of both micro- and macrovascular complications. Hyperglycaemia is associated with increased production of angiotensin II following RAAS over-activation in glomerular mesangial cells.¹¹⁶ Thus, mechanisms to block the RAAS are an important therapeutic target in people with type 2 diabetes and nephropathy.

Hypertension in people with type 2 diabetes

In nearly one-third of people with type 2 diabetes, hypertension is present at the time of their diagnosis. Hypertension and type 2 diabetes may be related to underlying diabetic nephropathy, to co-existing essential hypertension or to renovascular disease, or it may be part of the complex insulin resistance syndrome. Hypertension in people with type 2 diabetes is generally associated with expanded plasma volume, increased peripheral vascular resistance and low renin activity.¹¹⁷

The threshold for diagnosis and aims for hypertension control in people with type 2 diabetes vary according to national and international guidelines. In the UK, for the management of hypertension in people with diabetes and for those with a UACR of 70 mg/mmol (619.47 mg/g) or more and CKD, NICE guidance recommends a target blood pressure of less than 130/80 mmHg.³⁴ The American Diabetes Association and the American Society of Nephrology recommend a blood pressure of less than 140/90 mmHg in all patients with type 2 diabetes and renal disease.¹¹⁸ The Kidney Disease Improving Global Outcomes (KDIGO) guidelines recommend a blood pressure of less than or equal to 140/90 mmHg in those who have an albumin excretion rate (AER) of less than 30 mg per 24 hours (UACR >3 mg/mmol [>26.55 mg/g]), or less than or equal to 130/80 mmHg if the AER is greater than 30 mg per 24 hours (UACR >3 mg/mmol [>26.55 mg/g]) in people with type 2 diabetes.²³

The draft KDIGO 2020 guidance suggests a stricter control in all individuals with CKD targeting a systolic blood pressure <120 mmHg using a standardised blood pressure measurement technique, but acknowledges the lack of evidence in people who have diabetes and CKD.²³

There is little evidence base for recommending blood pressure targets in older people who have CKD. Most randomised controlled trials excluded people who were over 70 years of age (mean age 65 years: about 2.5% were older than 85 years of age) but some indirect inferences can be drawn from studies of older populations who do not specifically have CKD. While there is some evidence regarding the treatment of high blood pressure in much older people (that is, older than 80 years of age) from the Hypertension in the Very Elderly Trial (HYVET),¹¹⁹ it applies to a blood pressure target of 150/80 mmHg in people with CKD who have an eGFR >40 mL/min/1.73 m².

The Swedish Trial in Old Patients with Hypertension (STOP Hypertension), which compared antihypertensive treatments in preventing cardiovascular events in older people with diabetes (with a mean age of 75.8), supports an upright blood pressure target of no lower than 150/90 mmHg.¹²⁰

The KDIGO guidelines²³ suggest tailoring blood pressure treatment in older people with CKD to consider age, comorbidities and other therapies, with a gradual escalation of treatment and close attention to electrolyte disorders, acute deterioration in kidney function, orthostatic hypotension and side effects of medications.

Thus, it would seem reasonable to suggest a target upright systolic blood pressure of no less than 150 mmHg in people with diabetes and CKD aged over 75, taking into account side effects of medications and comorbidities.

The role of home and ambulatory blood pressure measurement

Although home and ambulatory blood pressure monitoring is thought to be more representative of real-life blood pressure, their exact role in the diagnosis and management of hypertension is unclear because a limited number of studies have been conducted in people with type 2 diabetes who have CKD. However, it is recognised that high ambulatory blood pressure measurement systolic pressures and nocturnal non-dipping are associated with increased mortality and a decline in eGFR.^{121–123} A small study of ambulatory blood pressure measurement in people with CKD, where 436 people who were hypertensive were prospectively followed up, showed that it was much more accurate in predicting both renal and cardiovascular outcomes than office blood pressure.¹²⁴ Self blood pressure monitoring and ambulatory blood pressure measurement utilises oscillometric assessment of blood pressure at the elbow, which may be influenced by irregularities of pulse and high pulse pressures. In the UK, NICE guidelines recommend confirming hypertension with 24-hour ambulatory monitoring (home BP monitoring where 24-hour BP monitoring is unsuitable or not tolerated) before starting or increasing antihypertensive agents.³⁴ However, there is no direct evidence supporting the use of either method for diagnosis of hypertension in people with type 2 diabetes and CKD.

Lifestyle modification and impact on blood pressure

There is good evidence from a number of observational studies and randomised controlled trials that salt intake, weight and body mass index (BMI), exercise frequency and alcohol intake all have a significant impact on blood pressure levels.^{35–38}

See box 1 on page 13 for more details.

Blood pressure lowering agents

In people with type 2 diabetes and CKD, three or more blood pressure agents are frequently required. There is increasing emphasis on individualisation of therapy. Eventually, the choice of agent is less important than the actual reduction in blood pressure that is achieved. There is little evidence to support the use of any particular agent in controlling blood pressure in CKD, nor are there any data to suggest the choice of second- or third-line medications. The exception to this rule is the use of ACEIs or ARBs in people with CKD who have proteinuria. ACEIs and ARBs have each been shown to be effective in delaying disease progression in people with type 2 diabetes who have microalbuminuria or established diabetic nephropathy. There is a need to escalate to the maximal doses of ACEI or ARB in people who have diabetes and albuminuria before moving on to additional agents in order to achieve the required blood pressure targets. However, there is no evidence that these agents are effective in the primary prevention of diabetic nephropathy. The use of ACEIs or ARBs in people with type 2 diabetes reduces microalbuminuria and retards the progressive loss of renal function.^{76,125–127} ARBs are said to provide renoprotection over and above their blood pressure lowering effect and short-term albuminuria reduction, and they are said to have a long-term favourable effect on renal prognosis.¹²⁸

Adherence with therapy

Non-adherence to antihypertensive treatment is common, with over 50% of people with apparent treatment resistance being non-adherent. This is especially so in those who are on multiple anti-hypertensive and other medications. Urine antihypertensive drug monitoring may help management of these individuals.¹²⁹ Therefore, it is important to assess adherence especially in those not achieving blood pressure control despite taking three or more antihypertensive agents in optimum doses.

RAAS blockade in people with type 2 diabetes without proteinuria

The use of RAAS blockade has significant benefits on cardiovascular and renal endpoints in people with diabetes, independent of their blood pressure lowering effect, as shown in the Heart Outcomes Prevention Evaluation (HOPE) trial and the European trial on reduction of cardiac events with perindopril in stable coronary artery disease.^{130,131} Whereas most guidelines favour the use of RAAS blockade as first-line treatment for people with diabetes, hypertension and CKD (the American Diabetes Association, the American Society of Hypertension, the International Society of Hypertension),^{132,133} the European Society of Cardiology / European Society of Hypertension guidelines from 2013 and the eighth Joint National Committee on Prevention, Detection, Evaluation and Treatment of High Blood Pressure from 2014 recommend the use of *any* class of antihypertensive agent in people with diabetes in the *absence* of proteinuria, but suggest the use of RAAS blockade as first-line treatment *only* in the presence of proteinuria.^{134,135}

This is based on the findings of 19 randomised controlled trials that enrolled 25,414 participants with diabetes, with a total of 95,910 patient years of follow-up. The results of this study from head-to-head randomised trials of RAAS blockade versus other antihypertensive agents failed to show superiority of RAAS blockade in people with diabetes and *no proteinuria*, and it suggested that any class of antihypertensive agents can be used in such individuals.¹³⁶

RAAS blockade in people with type 2 diabetes and proteinuria or microalbuminuria

In the UK, NICE guidance suggests offering a low-cost RAAS antagonist to people with CKD and diabetes who have: a UACR of 3 mg/mmol (26.55 mg/g) or more; hypertension and a UACR of 30 mg/mmol (265.49 mg/g) or more; or a UACR of 70 mg/mmol (619.47 mg/g) or more irrespective of hypertension or cardiovascular disease.¹³⁷ The favourable effects of RAAS blockade have been seen mainly in placebo controlled trials^{127,130} and it has been postulated that the benefits of RAAS blockade on renal outcomes was probably as a result of their blood pressure lowering effect.¹³⁸ Several major trials have also demonstrated clear benefits of ARB use in people who have diabetic nephropathy.^{139,140}

Use of dual blockade with ACEIs and ARBs in people with type 2 diabetic nephropathy

ACEIs and ARBs block the RAAS at different sites and, in theory, dual blockade should provide more effective and complete blockade of the RAAS. The rationale for dual blockade is based on a phenomenon called 'angiotensin II escape', whereby evidence suggests that standard doses of ACEIs only offer a partial blockade of the angiotensin-converting enzyme (ACE).¹⁴¹ It is said that enzymes such as chymase and cathepsin G can generate angiotensin II from angiotensinogen and other peptide substrates.¹⁴²

Several early studies had suggested that using a combination of ACEI/ARB provided additional benefit in diabetic nephropathy in terms of surrogate albuminuria lowering. However, there remains substantial controversy about whether ACEIs and ARBs should be combined, given that most of these studies were small in size and short in duration.

In one meta-analysis of 10 trials, 156 participants received a combination of ACEI/ARB and 159 received an ACEI only (the duration of the study was 8–12 weeks). The combination was shown to reduce proteinuria at the expense of statistically and clinically significant reductions in eGFR. There was a suggestion that this decrease could be secondary to a reduction in blood pressure alone.⁹² Most of the evidence base for combination dual blockade therapy initially came from studies about heart failure without any long-term data to support it (the candesartan and lisinopril microalbuminuria (CALM) study).¹⁴³ This study evaluated the effects of dual blockade of candesartan and lisinopril on blood pressure and microalbuminuria in 199 people with type 2 diabetes (the duration of the study was 24 weeks). At the end of the study, combination therapy was found to be significantly more effective in reducing UACR (50% with combination, 24% with candesartan and 39% with lisinopril) and diastolic blood pressure (16.3 mmHg, 10.4 mmHg and 10.7 mmHg reduction, respectively) than either agent alone. Criticisms of some of these studies were that there were no long-term follow-up data and that maximal doses of ACEIs were not used. It is also questionable whether the effects were specifically related to combination therapy or whether it was blood pressure reduction *per se* that was instrumental.

The ONTARGET study involved telmisartan and ramipril, and showed that the primary renal outcome (ie dialysis, doubling of serum creatinine and death) was similar for telmisartan (13.4%) and ramipril (13.5%), but was increased with combination therapy (14.5%, $p=0.037$). The combination therapy, although associated with reduced albuminuria, caused the greatest decline in eGFR.¹⁴⁴

The KDIGO guidelines provide specific advice on dual blockade.^{23,145} In the UK, a NICE guideline explicitly states that combination therapy should not be used. The European Renal Best Practice working group has the same viewpoint as NICE.¹⁴⁶ The Canadian Health Education Programme's (CHEP's) 2009 recommendation advised against the use of dual blockade for people with non-proteinuric CKD or in people with diabetes and normal urinary albumin levels.¹⁴⁷

Overall, therefore, there is no current evidence to suggest a beneficial effect of ACEI/ARB combination on the progression of diabetic nephropathy. Instead, combination therapy resulted in clinically significant decreases in eGFR and hyperkalaemia.¹⁴⁸ In one meta-analysis involving 17,337 people, the adverse effects of dual blockade revealed significantly high rates of discontinuation because of a worsening of renal function, hyperkalaemia and symptomatic hypotension.¹⁴⁹

It is to be noted, however, that most studies published so far with regard to dual blockade have involved people with *normal renal function* who did not have a clinically significant rise in serum potassium or creatinine with dual blockade. However, in real life, widespread use of these agents would most likely involve people with resistant hypertension with chronic renal impairment, and such individuals therefore will tend to have more of these side effects.

Aldosterone blockade in people with type 2 diabetic nephropathy

Aldosterone, the principal physiological mineralocorticoid, has deleterious effects on both the cardiovascular system and the kidneys. There is evidence to suggest that initial RAAS blockade suppresses aldosterone levels. However, due to the phenomenon of aldosterone

escape, aldosterone levels rise subsequently and can often exceed the baseline. ACEIs or ARBs do not directly block the effects of aldosterone at the receptor level.¹⁵⁰

Most evidence of the use of aldosterone antagonists like spironolactone, eplerenone and (more recently) finerenone come from heart failure trials. In one study of type 2 diabetes patient with early nephropathy and normal renal function, adding spironolactone to ACEI treatment was shown to be clinically useful and safe for people who showed aldosterone escape during ACEI treatment and who no longer showed maximal antiproteinuric effects of ACE inhibition.^{151,152} In another study of people with type 2 diabetes, macroalbuminuria and serum creatinine of less than 160 µmol/l, treatment with spironolactone was found to be superior to cilazapril in reducing albuminuria.¹⁵³ In that study, 50 mg of spironolactone was used and blood pressure of less than 135/85 mmHg was pre-treated with atenolol and hydrochlorothiazide before randomisation. The authors concluded that the superior effect of spironolactone was independent of its hypotensive effect, although 15% of people had to discontinue spironolactone because of hyperkalaemia.

There is also evidence for additive effects of eplerenone (an aldosterone antagonist that does not have the oestrogenic side effects of spironolactone) like in the other aldosterone antagonist trials. Eplerenone was found to have beneficial effects on microalbuminuria in people with type 2 diabetes when it was added to enalapril, although there was a much higher incidence of hyperkalaemia.¹⁵⁴ In the largest randomised controlled trial available using eplerenone, people with CKD, elevated urinary albumin levels and type 2 diabetes (177 people) received 50–100 mg of eplerenone and 91 people received a placebo. The addition of eplerenone to enalapril 20 mg per day resulted in a 40–50% reduction in AER by 12 weeks in the eplerenone group, but by less than 10% in the placebo group. Small reductions in eGFR and systolic blood pressure were noted, as was hyperkalaemia.¹⁵⁵

More recently, finerenone, a novel non-steroidal mineralocorticoid antagonist with greater receptor selectivity than spironolactone and eplerenone, has been shown to provide a greater reduction in proteinuria and end organ damage, compared with spironolactone or eplerenone. This was shown in the Mineralocorticoid Receptor Antagonist Tolerability – diabetic nephropathy (ARTS-DN) study involving 1,501 participants who were already receiving an ACEI or ARB (the mean age of the participants was 64.2 years, 37% had a UACR >30 mg/mmol [>265.49 mg/g] and 40% had an eGFR of 60 mL/min/1.73 m² or lower). Finerenone reduced the UACR at day 90 (relative to the baseline) more significantly than the placebo, and the pre-specified secondary outcome of hyperkalaemia leading to discontinuation was not observed either in the placebo or the finerenone group at various dosages. Also, there was no difference in terms of the incidence of a greater than 30% decrease of eGFR in either group. Thus 2.5–10 mg finerenone per day reduced albuminuria from the baseline in individuals with CKD and heart failure with a lower incidence of hyperkalaemia than spironolactone. It was postulated that this new mineralocorticoid receptor antagonist may be able to address the unmet medical need of safely managing albuminuria without effecting serum potassium in people with type 2 diabetes who have nephropathy. The strength of the study is that there was only a modest reduction in blood pressure at the highest dose of finerenone: quite unlike any other mineralocorticoid antagonist study in the past. The limitations of the study, however, include its short duration, the lack of a control group and that 60% of participants had an eGFR above 60 mL/min/1.73 m², which put them at relatively low risk of hyperkalaemia.¹⁵⁶

The FIDELIO-DKD study showed that finerenone, a non-steroidal selective mineralocorticoid receptor antagonist, lowered the risks of CKD progression and cardiovascular events in people with CKD and type 2 diabetes treated optimally with renin-angiotensin system blockade. Of the 5,734 people in the study, over 54% had eGFR <45 mL/min/1.73 m² and the median albumin creatinine ratio was 852 [IQR: 446–1634]. Finerenone, in doses of 10–20 mg

daily, was associated with improved renal (HR 0.82; 95% CI 0.73 to 0.93; NNT 29) and cardiovascular outcomes (HR 0.86; 95% CI 0.75 to 0.99; NNT 42) compared with placebo. In the study cohort, finerenone reduced systolic blood pressure by 3 mmHg and the incidence of hyperkalaemia of >6 mmol/L was 10% with discontinuation due to serious hyperkalaemia of 2.5%.^{157,158}

From the above evidence, it may be reasonable in the future to consider adding in a selective mineralocorticoid receptor antagonist in DKD patients with a serum potassium of less than 5 mmol with worsening albuminuria who are already on a maximal dose of ACEI or ARB.

Use of direct renin inhibitors in diabetic nephropathy

The use of aliskiren in people with type 2 diabetes and nephropathy has been shown to reduce AER, although no significant effects on renal outcomes have been noted. In the ALTITUDE trial, where aliskiren or a matching placebo was used on top of an ACEI or ARB in people with diabetic nephropathy, there were significant reductions in proteinuria but the trial was stopped early due to the low likelihood of ever demonstrating a benefit and the suggestion of an increased risk of non-fatal stroke, renal complications, hyperkalaemia and hypotension.¹⁵⁹ The drug has subsequently been withdrawn from the market.

In another trial involving 599 participants, aliskiren was used either alone or in combination with losartan for 6 months. This resulted in a reduction of UACR by 20% compared with the use of losartan alone. There were small differences in blood pressure between the two groups but no difference was found between the rates of adverse events.¹⁶⁰ Direct renin inhibitors are not currently recommended for use in diabetic nephropathy.

When should RAAS blockade be stopped?

Although ACEIs and ARBs are valuable blood pressure lowering agents in people with type 2 diabetes and CKD, they are not without their side effects.

a Hyperkalaemia

In the UK, NICE guidance suggests measuring serum potassium and eGFR before starting RAAS blockade and repeating the measurements 1–2 weeks after starting RAAS blockade and after each dose increase. NICE further says not to offer these agents if the person's pre-treatment serum potassium is greater than 5 mmol/L.³⁴ NICE guidance also suggests that these agents should be stopped if the serum potassium concentration increases to 6 mmol/L or more, and other drugs known to promote hyperkalaemia have been discontinued. However, recent NICE technology appraisals recommend the use of potassium binders, patiromer and sodium zirconium cyclosilicate, in outpatient care for people with persistent hyperkalaemia (≥ 6 mmol/L) and CKD stages 3b to 5 (non-dialysis), who are not taking an optimum dose of RAAS inhibitor because of hyperkalaemia.^{161,162}

b A drop in eGFR or an increase in serum creatinine

Given the basic pathophysiological mechanism of RAAS blockade, these agents cause a reduction in eGFR and urinary albumin excretion through efferent and afferent glomerular arteriolar dilatation, with a resultant fall in intra-glomerular blood pressure. A reversible reduction of eGFR of up to 30% can be expected. Greater reductions may indicate underlying renal artery stenosis.

NICE guidance states that if there is a decrease in eGFR of greater than 25% or an increase in serum creatinine of greater than 30% with RAAS blockade, renal function tests need to be repeated within 1–2 weeks. If the eGFR drops by 25% or more, or there is a change in serum

creatinine by 30% or more, NICE guidance suggests conducting further investigations to identify a cause of renal deterioration, such as volume depletion or non-steroidal inhibitor / potassium-sparing diuretic use. If no other cause for the deterioration in renal function is found, it is recommended to stop the RAAS blockade or reduce the dose to a previously tolerated lower dose, and add an alternative antihypertensive medication if required.^{137,163,164}

Consideration should also be taken where someone's baseline eGFR is already below 30, especially in those with congestive cardiac failure, where there may be a broader benefit terms of left ventricular function.

c Pregnancy

Given the potentially teratogenic nature of RAAS blockade drugs, the KDIGO guidelines suggest that the use of these drugs in women of childbearing age should be balanced with the risk of pregnancy.¹⁶⁵

d Inter-current illness

There are risks of large reductions in eGFR with RAAS blockade, particularly during intercurrent illness or with intravascular fluid depletion (diarrhoea, vomiting and high fever). It is therefore recommended to reduce the dose or to hold off ACEI or ARB use until recovery is made, because ensuing hypotension may cause an acute decline in eGFR in people with type 2 diabetes with CKD who are taking ACEIs or ARBs. It is recommended that people with type 2 diabetes should be advised to stop RAAS-blocking drugs during periods of acute illness and restarted 24–48 hours after recovery from the illness. These precautions should especially be taken if an individual is on a combination involving non-steroidal anti-inflammatory drugs or diuretics.^{166–169}

Other agents for blood pressure lowering in people with type 2 diabetes and nephropathy

Most of the evidence for the use of other antihypertensive agents (apart from ACEIs or ARBs) is extrapolated from the general population and there is little evidence of their specific use or rationale in people with type 2 diabetes and CKD.

a Calcium channel blockers

There is good evidence to suggest that non-dihydropyridine calcium channel blockers (verapamil and diltiazem) reduce proteinuria.^{87,169} A multicentre trial in people with type 2 diabetes and nephropathy suggested that adding a non-dihydropyridine calcium channel blocker to an ACEI-based regime can be effective at lowering residual albuminuria with or without a significant reduction in systolic blood pressure.¹⁷⁰ Thus non-dihydropyridine calcium channel blockers can be used as a valid additive or alternative to ACEIs or ARBs in people with type 2 diabetes, suggesting that their renal protective effects are over and above blood pressure lowering alone. Diltiazem and verapamil can induce bradycardia and heart block in combination with beta blockers; dihydropyridines such as amlodipine are more appropriate alongside beta blocker use.

b Beta blockers

Much of the bad publicity about beta blockers is related to the use of atenolol, which has been the most frequent comparator in most randomised controlled trials. However, beta blockers are not a homogenous class of drug, and agents like celiprolol, carvedilol and nebivolol have vasodilating properties and do not share the negative properties of atenolol

(that is, a lack of 24-hour antihypertensive effect and withdrawal effects). In the UK, NICE guidance does not favour beta blockers as the first-line choice in the treatment of hypertension in the general population. There is evidence that in people with type 2 diabetes with advanced CKD and a high risk of sudden death, beta blockers may prove to be beneficial by lowering heart rate apart from lowering sympathetic hyperactivity and preventing ventricular arrhythmias.^{171,172} A meta-analysis of beta blockers used to treat CKD supports the use of beta blockers in people with CKD who have heart failure, but it does not provide evidence of their efficacy in preventing mortality, cardiovascular events or renal disease progression in people with CKD who do not have heart failure.¹⁷³

c Diuretics

In the UK, NICE guidance prefers agents with a thiazide-like action such as chlorthalidone and indapamide, and this is relevant for individuals with CKD who have type 2 diabetes.³⁴ Chlorthalidone was used in the largest randomised controlled trial in hypertension (the Anti-hypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT) study).¹⁷⁴ The evidence base for indapamide is through the Hypertension in the Very Elderly Trial (HYVET).¹¹⁹ The PROGRESS trial involved a combination of indapamide and perindopril, and was shown to reduce the risk of stroke.¹⁷⁵ The additional advantage of indapamide is its potassium-depleting effect, and this may be convenient when it is combined with ACEIs or ARBs, particularly in those with type 2 diabetes who have CKD. Loop diuretics like furosemide would be particularly useful for treatment of hypertension in people with type 2 diabetes with advanced CKD stages 4–5, as fluid overload is invariably a major contributing factor in such individuals.

d Alpha blockers

Drugs like doxazosin could be an adjunctive treatment for hypertension in people with type 2 diabetes and CKD in whom other therapies have failed or not been tolerated, particularly if symptoms of prostatic hypertrophy are present. Alpha blockers are generally not recommended first line because of the common side effects of postural hypotension, tachycardia and headache.

e Centrally acting alpha adrenergic agonists

Centrally acting alpha adrenergic agonists cause vasodilation by reducing sympathetic outflow from the brain. Common agents in this category are methyldopa, clonidine and moxonidine. Doses of methyldopa and clonidine are not generally required to be reduced in people with CKD. Although moxonidine is extensively excreted by the kidney, one randomised controlled trial that compare it with a calcium channel blocker added to an ACEI or ARB plus a loop diuretic indicated that it is safe to be used in advanced CKD.¹⁷⁶ Common side effects of moxonidine include headache, tiredness, dizziness and gastrointestinal symptoms, which occur in 10–15% of people. These agents should not be used as a first-line treatment, but they are generally used in conjunction with other antihypertensive agents in people with type 2 diabetes who have hypertension.

Agents shown to have benefit in blood pressure reduction and outcomes but not currently licensed for this indication

f Endothelin A receptor antagonists

In the SONAR study (Study of Diabetic Nephropathy with Atrasentan), 4,711 adults with type 2 diabetes and kidney disease (eGFR of 25–75 mL/min/1.73 m², and a UACR between 33.9–565 mg/mmol [300–5,000 mg/g]), who had received stable treatment with renin–

angiotensin system blockade for at least 4 weeks were given oral atrasentan 0.75 mg/day. Atrasentan reduced primary composite renal endpoint of doubling of serum creatinine or ESKD, chronic dialysis for ≥ 90 days, kidney transplantation, or death from kidney failure (HR 0.65; 95% CI 0.49–0.88); SBP reduced in responders by -3.0 mmHg (-4.3 to -3.1) and in non-responders by -0.8 mmHg (-1.7 to -0.2).

These results suggest that endothelin receptor antagonists could play a role in preventing kidney failure in patients with DKD. Whether this is a direct effect of the class of drug or its modest BP lowering effect is difficult to postulate.¹⁷⁷

g Sodium glucose cotransporter-2 (SGLT-2) inhibitors

The CREDENCE (Canagliflozin and Renal Events in Diabetes with Established Nephropathy Clinical Evaluation) trial assessed the effects of the oral SGLT-2 inhibitor canagliflozin on renal outcomes in people with type 2 diabetes and albuminuric chronic kidney disease.^{178,179} Patients with DKD (eGFR between 30 to <90 mL/min/1.73 m² and UACR >33.9 –565 mg/mmol [>300 to 5,000 mg/g]) who had been treated with renin–angiotensin system blockade, were assigned to either canagliflozin 100 mg or placebo, and followed up for 2.62 years (median). The primary outcome was a composite of end-stage kidney disease (dialysis, transplantation, or a sustained eGFR of <15 mL/min/1.73 m²), a doubling of the serum creatinine level, or death from renal or cardiovascular causes. The risk of primary endpoint was reduced by 30% and that of end-stage kidney disease by 32%, along with significant reduction in the risk of cardiovascular events in the canagliflozin group compared with placebo.

Systolic blood pressure in the canagliflozin group was lower by 3.30 mmHg (95% CI, 2.73 to 3.87), and diastolic blood pressure by 0.95 mmHg (95% CI, 0.61 to 1.28). The observed cardiovascular and renal benefits happened despite very modest between-group differences in blood glucose level, weight, and blood pressure, raising the possibility of independent beneficial effect of canagliflozin.^{178,179}

3 Hypertension management and renin-angiotensin-aldosterone system blockade in people with type 2 diabetes, nephropathy and/or later stage CKD stages 4 and 5 (non-dialysis)

Recommendations

Identification and monitoring

- 1 We recommend regular monitoring of blood pressure, urine albumin, blood electrolytes and kidney function in people with diabetes and CKD stages 4 and 5 (Grade 1B).
- 2 We suggest, if blood pressure is uncontrolled, electrolytes are abnormal, or kidney disease is progressive they should be monitored 2 to 4 times a year depending on the stage of CKD and the individual's need (Grade 1B).

Target blood pressure

- 1 We recommend initiation of antihypertensive agents in people with diabetes and CKD stages 4 and 5, and UACR <3 mg/mmol (<26.55 mg/g) when blood pressure is >140/90 mmHg and aim for a target blood pressure of <140/90 mmHg during therapy (Grade 1B).
- 2 We suggest initiation of antihypertensive agents in people with diabetes and CKD stages 4 and 5 and UACR >3 mg/mmol (>26.55 mg/g) when blood pressure is >130/80 mmHg and aim for a target blood pressure <130/80 mmHg (Grade 2C).

Renin-angiotensin system blockade

- 1 We recommend the use of angiotensin converting enzyme inhibitor (ACEI) (or angiotensin receptor blocker (ARB) if ACEI is not tolerated) as the first-choice blood pressure lowering agent in people with diabetes and CKD stages 4 and 5 and micro/macroalbuminuria (Grade 1B).
- 2 We do not recommend the use of combinations of ACEIs and ARBs in people with diabetes and CKD stages 4 and 5 (Grade 2B).

Management of hyperkalaemia with renin-angiotensin system blockade

- 1 We suggest dietary advice, correction of acidosis and loop diuretic therapy to lower serum potassium as necessary in people with diabetes and CKD stages 4 and 5 for safe use of ACEI (or ARB) (not graded).
- 2 Consider the use of novel potassium binders in people with diabetes and CKD stages 3b to 5 (non-dialysis) if potassium is 6 mmol/L or higher, for continued and safe use of ACEi (or ARB) , or where people are not taking or are only taking sub maximal RAAS blockade because of hyperkalaemia (not graded).

Non-pharmacological management

- 1 We recommend dietary input to follow low sodium diet in all individuals with diabetes, advanced chronic kidney disease and high blood pressure (Grade 1B).

Audit standards

The following are suggested as audit standards for management of hypertension and RAAS blockade in people with diabetes and CKD stages 4 and 5:

- 1 Proportion of people with blood pressure <140/90 mmHg with diabetes and CKD stages 4 and 5 with UACR <3 mg/mmol (<26.55 mg/g).
- 2 Proportion of people with blood pressure <130/80 mmHg with diabetes and CKD stages 4 and 5 with UACR >3 mg/mmol (>26.55 mg/g).
- 3 Proportion of people who are taking an ACEI/ARB and have a serum potassium >5.5 mmol/L.
- 4 Proportion of people with potassium >5.5 mmol/L and being seen by dietitian.

Research recommendations

The following areas lack good quality evidence for management of hypertension in people with diabetes and CKD and hence further research may help:

- 1 To investigate the effect of intensive blood pressure lowering (<130/80 mmHg) on renal and cardiac outcomes in people with diabetes and CKD stages 4 and 5.
- 2 To investigate the impact of dual blockade with ACEI and ARB on renal and cardiac outcomes in people with diabetes and CKD and proteinuria
- 3 To investigate the effect of long-term use of novel potassium binders together with renal angiotensin system blockade on renal and cardiac outcomes in people with diabetes and CKD stages 4 and 5.
- 4 To investigate the impact of new agents, namely endothelin A receptor antagonists and SGLT-2 inhibitors, on blood pressure in people with diabetes and CKD stages 4 and 5.

Introduction

Advanced stages of CKD, particularly stages 4 and 5, are associated with hyperkalaemia, fluid retention and anaemia requiring erythropoiesis stimulating agents which may further increase blood pressure. Hyperkalaemia (>5.5 mmol/L) is present in 31% of people in advanced kidney disease clinics.¹⁸⁰ In a large population, hyperkalaemia (>5.5 mmol/L) occurred more frequently in people with CKD (eGFR <60 mL/min/1.73 m²) on renin-angiotensin blockers than without (8 vs. 2 events per 100 patient-months; p<0.001).¹⁸¹ Hyperkalaemia is more common in people with CKD and diabetes than in those with CKD without diabetes.¹⁸² In a blood pressure control trial in people with CKD the risk of hyperkalaemia was seven times higher in patients with eGFR <30 mL/min/1.73 m² compared with eGFR >50 mL/min/1.73 m² and seven times higher with ramipril compared with amlodipine.¹⁸³ Hence BP control in people with diabetes and CKD stages 3–5 particularly with an ACEI or ARB requires careful monitoring and management of serum potassium. The prevalence of primary aldosteronism in people with diabetes and resistant hypertension is 14%, and this should be considered when BP is difficult to control.¹⁸⁴

Identification and monitoring of people with diabetes and CKD stages 4 and 5

The rise in blood pressure in people with diabetic nephropathy is associated with higher mortality and increased risk of macro and micro vascular complications;^{185,3} and treatment lowers cardiovascular events, strokes and all-cause mortality.^{186–188} Hence people with diabetes and CKD stages 4–5 should be regularly screened to identify and manage high blood pressure. It is necessary to identify those with hypertension early to avoid delays in treatment; while avoiding unnecessary anxiety and the inconvenience related to frequent visits to doctors and nurses. With the use of RAAS blockers, monitoring of serum potassium is important to avoid dangerous hyperkalaemia.¹⁸⁹ Frequent blood testing will also identify people who are more likely to progress to renal replacement therapy.¹⁹⁰ Most clinical trials have monitored individuals' clinical characteristics and laboratory values every 3 to 12 months and have demonstrated identification of new onset hypertension, proteinuria and hyperkalaemia in this time frame.^{183,191,192} The recommendations for monitoring by NICE are: twice a year for CKD stage 3a, ≥ 2 times a year for CKD stage 3b, 3 times a year for CKD stages 3 and 4 and ≥ 4 times a year for CKD stage 5. However, this can be tailored according to the individual's needs.¹³⁷ During the first consultation, ambulatory (or home) blood pressure monitoring should be offered to confirm the diagnosis of hypertension.³⁴ Measurement of sitting and standing blood pressure may be useful to diagnose postural hypotension which may contribute to symptoms and standing blood pressure may be a better target. Blood pressure should be measured by standardised technique in a quiet, comfortable environment, on an outstretched supported arm, using a properly calibrated machine with an appropriate cuff [as suggested by the BIHS](#).

Target blood pressure in people with diabetes and CKD stages 4 and 5, with or without significant albuminuria (UACR >3 mg/mmol [>26.55 mg/g])

Several observational and prospective studies have demonstrated the significant impact of blood pressure on mortality, cardiovascular events and renal failure in people with diabetes and CKD.³ Among those from advanced CKD clinics, high blood pressure (particularly systolic) is associated with progression to dialysis and mortality.¹⁹³

Very few studies have examined the impact of tight blood pressure control in people with diabetes and CKD stages 4 and 5. Some studies have examined the role of intensive blood pressure lowering in people with diabetes and mild CKD; a small proportion of people demonstrating the advantage of lowering blood pressure below 140/90 mmHg but not below 130/80 mmHg.¹⁹⁴ In a study of African-American people with non-diabetes CKD and eGFR 20–65 mL/min the tight blood pressure (achieved 128/78 mmHg) control arm suffered similar renal endpoints compared to less tight blood pressure (achieved 141/85 mmHg) control arm.¹⁹⁵ In the SPRINT trial, which included 2,646 non-diabetic patients with eGFR 20–60 mL/min/1.73 m², intensive blood pressure control (target <120 mmHg) was not associated with improved composite renal outcomes, compared with standard control in patients with chronic kidney disease (target <140 mmHg).¹⁹² However, no participants with diabetes were included in the SPRINT trial. In another large randomised controlled trial of high risk individuals with diabetes (ACCORD-BP), the intensive blood pressure control arm (target SBP <120 mmHg; achieved 119 mmHg) was associated with a higher chance of having a eGFR <30 mL/min/1.73 m² (99 vs 52 events; $p<0.001$) than the normal blood pressure control arm (target SBP <140 mmHg; achieved 133 mmHg), without any benefit in reducing cardiovascular complications.¹⁹¹ This trial excluded people with creatinine above 1.5 mg/dL (approximate eGFR 50 mL/min/1.73 m²), and mean creatinine at baseline was 0.9 mg/dL

(approximate eGFR 91 mL/min/1.73 m²). In the same study there was no difference in new onset microvascular complications with intensive blood pressure control and half of the people who had progressive renal disease did not have albuminuria.¹⁹⁶ However, a pooled analysis of SPRINT and ACCORD-BP with 14,094 people followed for 3.26 years suggests a 18% risk reduction of cardiovascular events and cardiovascular deaths.¹⁹⁷ However, lowering blood pressure <130/80 mmHg may be associated with unwanted side effects and individuals should be involved in the decision-making process.¹⁹⁸

Several studies have shown that presence of significant albuminuria is associated with poor cardiovascular outcomes and reduction of albuminuria is associated improvement.¹⁹⁹ Analysis of data from the RENAAL study, a trial of ARB in diabetic nephropathy, demonstrated an approximate doubling of risk of cardiovascular outcome with high albuminuria at baseline (ACR >3g/g compared with <1.5 g/g of creatinine), and 18% lowering of the cardiovascular events with 50% lowering of albuminuria.²⁰⁰ The evidence for better outcome with tighter blood pressure control with high albuminuria is mainly observational and derived from post hoc analysis of large randomised controlled trials. The RENAAL study which included a significant number of people with CKD stages 3 and 4 demonstrated baseline higher risk with SBP >140 mmHg (no difference between <130 compared with 130–140 mmHg), and a 23% risk reduction for ESKD with achieved BP <140/90 compared with >140/90.²⁰¹ Analysis of data from two large ARB trials (IDNT and RENAAL) indicate that the benefits of cardiovascular risk reduction exist with SBP <130 mmHg (particularly when albuminuria was reduced to lower levels). However, the risk increased with SBP <120 mmHg.²⁰² Post-hoc analysis of IDNT trial demonstrated a benefit in reduction of heart failure events with SBP <130 mmHg but possible increased risk with SBP <120 mmHg; and a DBP <85 mmHg was associated with increased risk of MI and CHF.²⁰³ Hence there is a suggestion of better cardiovascular outcomes with reduction of systolic blood pressure below 130 mmHg but not below 120 mmHg. With target blood pressure <130/80 mmHg in the STENO 2 randomised trial there was reduction in cardiovascular mortality; however all most patients had CKD stages 1 and 2 and in the presence of other interventions it is difficult to tease out the effect of tight BP control.²⁰⁴ Thus a lower target for blood pressure <130/80 mmHg may be suggested in patients with significant albuminuria as suggested by other guidelines (KDIGO/NICE) but stronger evidence is needed.^{205,137} The draft KDIGO 2020 guidelines suggest a lower blood pressure target <120 mmHg systolic in all patients with CKD with diabetes irrespective of the degree of proteinuria, while acknowledging that the degree of evidence in support of tighter control is low (evidence grade 2B), the clinical risk of adverse events with low blood pressure target, particularly when measured in non-standardised manner is high; thus allowing clinicians to target higher blood pressure when necessary.²³ This guideline proposes a target blood pressure of <140/90 mmHg for all individuals with diabetes, no significant proteinuria and CKD stages 4 and 5, as improved cardiovascular outcomes have been demonstrated in randomised controlled trials with blood pressures <140/90, but inconsistent results with lower targets. Whereas the proposed target is <130/80 mmHg is for people with significant proteinuria as it is associated with reduction in proteinuria in diabetic kidney disease which may improve renal and cardiovascular outcomes.

Renin-angiotensin system blockade for blood pressure control in people with diabetes and CKD stages 4 and 5

In a study of African-American people without diabetes and with eGFR 20–65 mL/min/1.73 m², use of the ACEI ramipril was associated with significant reduction in clinical composite outcome compared to metoprolol (22% [95% CI, 1%–38%; p=0.04]) or amlodipine (38% [95% CI, 14%–56%; p=0.004]).¹⁹⁵ In a randomised controlled trial of people with diabetes (30%

with mild CKD) use of enalapril was associated with fewer cardiovascular events compared with nisoldipine (5 vs. 25; $p < 0.001$).²⁰⁶ In a recent meta-analysis of 119 trials, use of ACEI or ARBs in 64,768 participants with CKD was associated with reduced risk of kidney failure compared with other antihypertensives (odds ratios of 0.65 [95% CI 0.51–0.80] for ACEIs and 0.75 [95% CI 0.54–0.97] for ARBs).²⁰⁷ Hence ACEIs should be used in people with diabetes and CKD stages 4 and 5, with careful monitoring of kidney function and serum potassium. In a meta-analysis treatment with ACEIs in people with diabetes was shown to reduce all-cause mortality, cardiovascular mortality and cardiovascular events but not with ARBs.^{208,209} A recent network meta-analysis showed reduction in ESKD with ACEI and/or ARB but did not demonstrate overall survival benefit.²¹⁰ Thus there is strong evidence for use of ACEI/ARB as the first choice antihypertensive in people with diabetes and CKD with eGFR >30 mL/min/1.73 m². Participants in large randomised controlled trials of people with diabetes had eGFR no lower than approximately 25 mL/min/1.73 m² in REENAL and 38 mL/min/1.73 m² in micro HOPE.^{201,211} Though hyperkalaemia and rapid decline in kidney function is an issue in people with advanced CKD, recent analysis of 3,909 individuals with CKD stages 4–5 suggested continuing treatment with ACEI or ARB was associated with cardiovascular benefit. The impact of withdrawal of ACEI or ARB on progression of CKD is being investigated by the STOP ACE randomised controlled trial.^{212,213}

In a study of combination therapy of ACEI with ARB in patients who have diabetes with UACR >33.9 mg/g (>300 mg/g) and eGFR $30–90$ mL/min/1.73 m², there was no difference in mortality but a significant increase in hyperkalaemia (6.3 events vs. 2.6 events per 100 person-years with monotherapy; $p < 0.001$) and acute kidney injury (6.7 vs. 0.2 events per 100 person-years, $p < 0.001$).⁹⁵ Combination of ACEI with ARB was not associated with benefit in primary endpoints but more side effects hence should be avoided. However, a network meta-analysis suggested a potential benefit of dual blockade if it can be administered safely, hence the need for further trials of dual-blockade in people who have diabetes with CKD and albuminuria.^{210,145}

A rise in serum creatinine up to 30% is not uncommon and rise of potassium by 0.5 mmol/L is not uncommon with initiation of ACEI therapy.¹⁰⁰ A post-hoc analysis of the ACCORD-BP trial demonstrates $>30\%$ rise in creatinine identifying people at risk for cardiovascular and all-cause mortality, but only associated with adverse renal outcome in the standard arm and not intensive therapy arm.¹⁹⁷ Thus a mild rise in creatinine may not require any change in planned therapy. Hence no modification of ACEI or ARB therapy is necessary if the rise in creatinine from baseline is $<30\%$ or drop in eGFR is $<25\%$.

Addition of spironolactone and further inhibition of the renin-angiotensin-aldosterone system may provide additional anti-proteinuric effect as seen in small studies and merits further large trials with more clinically relevant outcomes.²¹⁴



Risk of AKI

During an acute illness particularly with volume depletion, there is a risk of acute kidney injury (AKI) in patients with diabetes and CKD stages 4 and 5 who are on ACEI or ARB therapy. Such risk can be avoided by stopping the ACEI or ARB for the duration of the illness and 24–48 hours post recovery.

Management of hyperkalaemia with renin-angiotensin system blockade in people with diabetes and CKD stages 4 and 5

Hyperkalaemia is common in people with diabetes and CKD. It is very common ($>30\%$) in advanced CKD patients managed in the low-kidney-clearance clinics.¹⁸⁰ The cause of such hyperkalaemia can be multifactorial; including renal failure, type 4 renal tubular acidosis,

diet and drugs. The presence of hyperkalaemia limits the use of renin-angiotensin-axis inhibitors. The chronically high potassium levels have been traditionally controlled with restricted diet, diuretics and avoiding drugs that cause hyperkalaemia.

Traditionally hyperkalaemia has been managed with dietary potassium restriction and correction of acidosis, if present with bicarbonate therapy. However, the new potassium binding agents have been tested for safety and efficacy in randomised controlled trials for management of chronic hyperkalaemia in CKD patients. They cause an early and sustained lowering of potassium in people with CKD on RAAS blocker therapy.²¹⁵ In 306 individuals with diabetes and CKD stages 3 to 4, treated with RAAS blockade (ACEI/ARB ± spironolactone), use of a novel potassium binding polymer (patiomer) was associated with significant and sustained decrease in serum potassium over 52 weeks.²¹⁶ In a study of 237 CKD patients the same potassium binder was able to reduce serum potassium by 1 mmol/L over 4 weeks.²¹⁷ In another study of 243 participants over >50% of whom had diabetes, the potassium binder achieved approximately 1 mmol/L reduction in serum potassium over 4 weeks in individuals with and without heart failure.²¹⁸ The treatment with patiomer was associated with decrease aldosterone levels and decreased blood pressure which may provide additional benefits.²¹⁹ However, the above-mentioned trials are of short duration and the possible ACEI or ARB use facilitated with potassium binders, has not been shown to improve cardiovascular events or mortality. However, in the recent AMBER study, in people with resistant hypertension and chronic kidney disease (50% with diabetes), patiomer enabled more patients to continue treatment with spironolactone with less hyperkalaemia.^{220,221}

Non-pharmacological management of hypertension in people with diabetes and CKD stages 4 and 5

In randomised controlled trials dietary sodium restriction in people with CKD is associated with significant lowering of blood pressure, but longer-term benefits of dietary intervention are unknown.^{222,223} The dietary advice is best provided by a trained dietitian due to the complex and frequently changing needs in this group of people. Individuals with CKD stages 4 and 5 would benefit most from this and are best managed in a multidisciplinary clinic with expert nurses and dietitians. Dietary potassium restriction is useful but clinical trial evidence is yet to be generated. Regular exercise tends to improve quality of life, eGFR decline, HBA1C, BMI without any adverse effects in people with diabetes and CKD stages 3–5.²²⁴ People should be also advised to quit smoking which is known to improve blood pressure in hypertensive individuals.

4 Hypertension management and renin-angiotensin-aldosterone system blockade in people with diabetes and CKD stage 5 on dialysis (5D)

Recommendations

The following are recommendations for blood pressure control in people with diabetes and CKD stage 5D.

- 1 We recommend that ambulatory blood pressure measurement or home blood pressure measurement should be used to monitor blood pressure in people with diabetes who are on dialysis (Grade 1C).
- 2 Where ambulatory blood pressure measurement or home measurement are not feasible to monitor blood pressure in people with diabetes who are on dialysis, we suggest using pre-, intra- and post-dialysis standardised blood pressure measurements for people who are on haemodialysis, and using standardised clinic blood pressure measurements for people who are on peritoneal dialysis (Grade 2D).
- 3 We recommend volume control as first-line management to optimise blood pressure control in people with diabetes who are on dialysis (Grade 1B).
- 4 We suggest salt restriction to less than 5 g per day to optimise blood pressure control in people with diabetes who are on dialysis (Grade 2C).
- 5 We suggest a target upright interdialytic blood pressure of <140/90 mmHg for people with diabetes who are on dialysis. Individualisation of the blood pressure target may be indicated in other people who are burdened with multiple comorbidities, in order to reduce adverse events of blood pressure lowering (Grade 2D).
- 6 We recommend that intradialytic hypotension should be avoided in people with diabetes who are on haemodialysis (Grade 1B).
- 7 We suggest using ACEIs or ARBs (but not in combination), beta blockers and calcium channel blockers to reduce cardiovascular complications in people with diabetes and hypertension who are on dialysis (Grade 2B).
- 8 We suggest the use of diuretics in people with diabetes who are on dialysis and have residual kidney function (Grade 2C).

Audit standards

The following are suggested as audit standards for blood pressure control in people who have diabetes and CKD stage 5D.

- 1 The proportion of people with diabetes receiving haemodialysis who achieve an upright interdialytic blood pressure target of <140/90 mmHg.
- 2 The proportion of people with diabetes receiving haemodialysis who achieve an upright interdialytic blood pressure target of <140/90 mmHg without the use of blood pressure lowering medication.
- 3 The proportion of people with diabetes receiving haemodialysis with an interdialytic blood pressure of >140/90 mmHg who are not being treated with ACEIs, ARBs, beta blockers or calcium channel blockers in whom such drugs are not contraindicated.

- 4 The proportion of people with diabetes receiving dialysis who are on diuretics following commencement of dialysis.

Areas that require further research

The following areas lack good-quality evidence for blood pressure control in people with diabetes and CKD stage 5D, and hence further research is necessary.

- 1 Which blood pressure measurement should be used to predict left ventricular hypertrophy (LVH) and mortality in people with diabetes who are on dialysis: pre-dialysis, post-dialysis, home or ambulatory blood pressure measurement?
- 2 What is the optimal upright blood pressure target for people with diabetes who are on dialysis?
- 3 Can bioimpedance spectroscopy devices be used to determine a target weight and predict the risk of cardiovascular morbidity for people with diabetes who are on dialysis?
- 4 Does treatment with ACEIs, ARBs, beta blockers or calcium channel blockers lower blood pressure in people with diabetes who are on dialysis reduce cardiovascular morbidity and mortality?
- 5 Is there a role for diuretic therapy in people with diabetes who are on dialysis and have residual kidney function?
- 6 Does salt restriction (to less than 5 g per day versus less than 6 g per day) in people with diabetes who are on dialysis influence blood pressure control or cardiovascular outcome?

Introduction

Hypertension is a common finding in people with diabetes as well as those with CKD stage 5D. Elevated blood pressure,^{225,226} diabetes^{3,227} and CKD^{228–230} are all major risk factors for adverse cardiovascular events.

According to the UK Renal Registry and the European Renal Association – European Dialysis and Transplant Association (ERA-EDTA) Renal Registry, 23–36% of incident dialysis patients had diabetes as their primary renal disease.^{231,232} People with diabetes who are on haemodialysis have a poorer survival rate compared with dialysis patients who do not have diabetes.^{233,234} This is mainly due to cardiovascular disease.^{235–237} Control of hypertension in hypertensive dialysis patients was shown to be associated with improved survival.²³⁸

It is therefore logical that, in order to reduce cardiovascular risk and improve survival, optimal blood pressure control should be achieved in people with diabetes and CKD stage 5D. However, there is insufficient evidence from data in the published literature to decide how best to manage blood pressure in people with diabetes who are on dialysis. This is in part because people with CKD, including those with stage 5D, are ‘often’ excluded from clinical trials of hypertension.

There are emerging, although not consistent, data delineating how best to measure blood pressure, to target blood pressure and to use pharmacological and non-pharmacological therapies to optimise blood pressure control in people with CKD stage 5D. However, these data are not specific to the population with diabetes.

Furthermore, blood pressure control in people who are on dialysis is complex. Many factors affect blood pressure in people who are on dialysis, including fluid status, salt intake, sympathetic nervous system activity and the renin-angiotensin-aldosterone system (RAAS). People with diabetes who are undergoing haemodialysis often have autonomic

dysfunction,²³⁹ which increases the risk of cardiovascular instabilities during dialysis. Haemodialysis causes severe orthostatic reduction in cerebral blood flow velocity in people with diabetes and may subsequently increase the risk of cerebrovascular injury post haemodialysis.²⁴⁰ This makes management of hypertension in people with diabetes who are on dialysis even more challenging.

Blood pressure measurement in people with diabetes who are on haemodialysis

In UK dialysis units, measuring pre- and post-dialysis blood pressure is the standard technique for monitoring blood pressure in people who are on dialysis. However, blood pressure measurement in people who are on haemodialysis is complex. There are conflicting data as to whether blood pressure measurements pre- and post-dialysis are predictive of interdialytic blood pressure in comparison with ambulatory blood pressure measurement and/or home blood pressure measurement. Ambulatory blood pressure measurement is considered to be the most accurate method for studying blood pressure in people who are on haemodialysis²⁴¹ and in the general population it provides a more accurate prediction of cardiovascular outcomes in comparison with clinic blood pressure measurement.²⁴² A meta-analysis of 18 studies that involved 692 people who were on dialysis showed that pre-dialysis blood pressure and post-dialysis blood pressure are imprecise estimates of interdialytic ambulatory blood pressure.²⁴³ People with diabetes were included in most of these studies^{235,241} at a rate that varied from 8% to 54%. The presence of diabetes made no difference to the outcome.

In an extensive review of the literature by Agarwal *et al*,²⁴⁴ evidence from several studies was presented to show that, in people on haemodialysis, blood pressure measurement at home²⁴⁵ or ambulatory blood pressure measurement^{246,247} are stronger predictors of LVH²⁴⁸ and mortality^{249,250} compared with blood pressure obtained in the dialysis unit. In predicting LVH, weekly average home systolic blood pressure measurement was similar to interdialytic ambulatory blood pressure measurement and was superior to pre-dialysis and post-dialysis blood pressure measurement.²⁴⁸ In contrast to home blood pressure measurement, ambulatory blood pressure measurement can diagnose nocturnal non-dipping and offers great insights into circadian rhythm.²⁵¹ Loss of diurnal rhythm, which is a feature of diabetic nephropathy, is reported to lead to worse outcomes in people who are on dialysis.²⁴⁶ In a study of 89 people who are on haemodialysis by Liu *et al*, the incidence of cardiovascular events and deaths were 3.5–9 times higher in non-dippers (that is, those who lose their diurnal blood pressure variation) compared with dippers.²⁵² Ambulatory blood pressure measurement can therefore be advantageous in selecting high-risk individuals and can guide treatment. However, to date, there have been no specific studies to address ambulatory blood pressure measurement in people with diabetes who are on dialysis.

While 24-hour ambulatory blood pressure measurement (ABPM) is considered gold standard in predicting outcomes, it is resource intensive, impractical for long-term monitoring of BP control and often not tolerated by patients. Interdialytic home blood pressure (HBPM) is close to ABPM in predicting outcomes, but has high attrition rate for long-term monitoring. As routinely collected peri-dialytic BP measurements guide interventions in the majority of dialysis units, it is essential to ensure routinely collected dialysis unit BP readings are measured in a standardised manner in accordance with recommended guidelines.

Target blood pressure in people with diabetes who are on dialysis

The relationship between blood pressure level and cardiovascular outcome is unclear in individuals who are on dialysis. Observational studies have shown an increased risk of mortality in people who are on haemodialysis who have a low pre- or post-dialysis systolic blood pressure of less than 110 mmHg,²⁵³ and in those who are on peritoneal dialysis with a pre-dialysis systolic blood pressure of <110 mmHg.²⁵⁴ Further observational studies in haemodialysis cohorts^{255,256} continued to show a reverse epidemiology phenomenon, with the highest mortality rate being in groups with lower pre-dialysis blood pressures. Recent observational relationship studies using peri-dialytic measurements have consistently shown a 'U'- or 'J'-shaped relationship with mortality. For example, analysis of Dialysis Outcomes and Practice Patterns Study (DOPPS) data, at both facility and individual patient level, found lowest mortality in those with a pre-HD SBP of 130 to 159 mmHg (facility level) and <130 mmHg (individual patient level); the facility-level analysis compensates for unmeasured confounding, albeit in an imperfect manner.²⁵⁷ Similarly, the CRIC Investigators reported a pre-HD SBP of 138 to 166 mmHg to be associated with lowest risk of cardiovascular events.²⁵⁸

Interestingly, in the Tassin group in France where the 5-year survival rate of 87% is the best reported in people who are on haemodialysis, the pre-dialysis blood pressure that was achieved was less than 130/85 mmHg (mean arterial pressure (MAP) <101 mmHg).²⁵⁹

Prospective randomised controlled studies on the effect of ARBs,²⁶⁰ ACEIs,²⁶¹ beta blockers²⁶² and calcium channels blockers²⁶³ on cardiovascular events have been conducted to evaluate the roles of these agents in people who are on dialysis.

Two meta-analyses have shown that blood pressure treatment in patients who are on dialysis is associated with improved outcome. The first analysis was by Heerspink *et al*, published in 2009.²⁶⁴ This meta-analysis included eight randomised trials that provided data from 1,679 people who are on dialysis, of whom 588 had diabetes. The trials included people who were on haemodialysis and peritoneal dialysis. The analysis showed that blood pressure lowering treatment was associated with lower risks of cardiovascular events, all causes of mortality and cardiovascular mortality, and that the effect seemed to be consistent across a range of groups that were included in the studies. Reduction in systolic blood pressure was similar, regardless of whether the person had diabetes or antihypertensive drug use. Similarly, the second meta-analysis by Agarwal and Sinha (also published in 2009)²⁶⁵ showed that in people with hypertension who were on haemodialysis, antihypertensive therapy reduced the combined hazard ratio for cardiovascular events by 31–38% compared with the placebo group. The meta-analysis showed that blood pressure lowering was well tolerated, with no suggestion of increased adverse events in people with diabetes. The analysis showed no difference in cardiovascular outcomes caused by different drug classes and the data from the two meta-analyses suggest that RAAS blockers, beta blockers and calcium channel blockers are all suitable for use in people who are on dialysis.

However, a randomised controlled trial by Agarwal *et al* in 2014,²⁶⁶ including 200 people who were on haemodialysis of whom nearly half had diabetes, showed that a beta blocker-based hypertensive treatment was superior to an ACEI-based treatment in preventing cardiovascular morbidity in those who are on dialysis.

Irrespective of the type of the antihypertensive agents that are used, the timing of the administration of such agents in relation to dialysis treatment needs to be taken into account when prescribing antihypertensive drugs for people who are on dialysis. ARBs, calcium channel blockers and alpha-blockers are not cleared with dialysis. However, ACEIs (apart from fosinopril) and a number of beta blockers are largely cleared on dialysis.²⁶⁷

To date, optimum blood pressure goals for patients who are on dialysis (including people with diabetes) have not been defined in randomised prospective controlled trials.²⁶⁷ The Kidney Disease Outcomes Quality Initiative (KDOQI) recommends a pre-dialysis blood pressure goal of less than 140/90 mmHg and a post-dialysis blood pressure goal of less than 130/80 mmHg.²⁶⁸ However, this is largely based on studies that were performed in the non-dialysis population who have normal renal function. People who are on haemodialysis have different characteristics to the general population. For example, studies have shown that people who are on dialysis lose their diurnal blood pressure variation (that is, they are non-dippers), which is an independent risk factor for LVH and subsequent adverse cardiovascular outcome.²⁴⁶ People who are on haemodialysis also have increased pulse pressure, which is associated with adverse cardiovascular outcome.²⁶⁹ Therefore, the KDOQI-recommended blood pressure target may not be applicable to the haemodialysis population.

Furthermore, people with diabetes who are on dialysis are at increased risk of haemodynamic instabilities and orthostatic intolerance post-dialysis, and therefore a blood pressure that is higher than 140 mmHg systolic may be indicated in the presence of significant orthostatic change in blood pressure. A randomised controlled trial is needed to identify the optimal blood pressure target for people with diabetes who are on dialysis.

As for patients who are on peritoneal dialysis, International Society of Peritoneal Dialysis (ISPD) guidelines recommend a target of <140/90 mmHg for self-measured home blood pressure readings.²⁷⁰ However, this is based on a number of small observational studies and not specifically for those who have diabetes. There is no randomised controlled trial evidence is currently available.

Volume control in people with diabetes who are on dialysis

Increased extracellular volume or volume overload is an important contributor in the pathogenesis of high blood pressure in people who are on dialysis.²⁷¹ Removal of extracellular volume without causing intolerable hypotension defines the 'dry weight'²⁷² that was first reported by Thomson in 1967.²⁷³ This is difficult to define clinically. Achieving dry weight and normalising blood pressure is not immediate and can take months, which is something that is best described as a 'lag phenomenon'.²⁷⁴

In the Tassin group, hypertension control without medication, achieved by aggressive control of extracellular volume and dietary sodium intake, was shown to be the best single marker of survival in 449 people who were on haemodialysis who were followed for 20 years.²⁵⁹

Observational studies showed that volume control is associated with improvement in blood pressure in the majority of people on haemodialysis²⁷⁵ and peritoneal dialysis.²⁷⁶ A randomised controlled trial (DRIP) showed that volume control in haemodialysis improves blood pressure control.²⁷⁷ In that study, 150 people were randomised to an additional ultrafiltration group (40/100 had diabetes) or control group (19/50 had diabetes). Without increasing time or frequency of haemodialysis, reduction in dry weight (defined by clinical signs and symptoms) resulted in a reduction in interdialytic ambulatory blood pressure, leading to the conclusion that dry weight reduction is an effective strategy in blood pressure control in people who are on haemodialysis.

The concept that 'volume control' improves blood pressure control is further supported by the increasing reports that daily dialysis^{278,279} or nocturnal dialysis^{280,281} improves blood pressure and reduces LVH with less risk of inducing intradialytic hypotension. Reducing the risk of intradialytic hypotension is important. An observational study by Shoji T *et al* showed that haemodialysis-associated hypotension is an independent risk factor for 2-year mortality in people who are on haemodialysis.²⁸²

The risk of intradialytic hypotension increases with an ultrafiltration rate of greater than 10 ml/kg/hr and was reported in the Dialysis Outcomes and Practice Patterns Study (DOPPS) (which included 16,420 patients on haemodialysis) as an independent risk factor for mortality.²⁸³ This is similar to another study with 5 years' follow-up data by Movilli,²⁸⁴ in which an ultrafiltration rate of over 12.7 ml/kg/hr was identified as an independent risk factor for mortality given the risk of hypotension-related serious adverse events especially in people who are on dialysis and who have diabetes; however, this strategy requires close supervision and markers to assess volume status.

Bioimpedance spectroscopy devices,²⁸⁵ brain natriuretic peptide (BNP) level,²⁸⁶ assessment of vena cava diameter²⁸⁷ and ultrasound lung water measurement²⁸⁸ have been used to determine dry weight. Of these, bioimpedance spectroscopy is most widely studied.²⁸⁹ Further studies are needed to explore and evaluate the role of bioimpedance spectroscopy devices as markers of volume status, especially in people with diabetes who are on dialysis.

Salt restriction in people with diabetes who are on dialysis

Reducing dietary salt to control blood pressure in people who are on dialysis was first reported by Hegstrom RM *et al* in 1961.²⁹⁰ Salt restriction to 1 g per day or less helps to decrease thirst and to control interdialytic weight gain in people who are on haemodialysis.²⁹¹

Evidence for the association between salt restriction and blood pressure control in people who are on dialysis comes from observational studies where dietary salt restriction was in combination with strict volume control. Craswell *et al*²⁹² (in a study of 89 patients who were on dialysis), Covic *et al*²⁹³ (in a study of 286 people) and Ozkahya *et al*²⁹⁴ (in a study of 218 people) all showed that salt restriction to less than 5 g per day along with strict volume control led to a significant reduction of blood pressure and interdialytic weight gain. Similarly, in the Tassin group, dietary salt reduction to <5 g per day along with extracellular volume control was shown to normalise blood pressure in people who were on haemodialysis.²⁵⁹

In a cross-sectional study by Kayikcioglu *et al* in 204 people on dialysis, dietary salt restriction to 5 g per day along with dialysate sodium reduction, led to a reduction in interdialytic weight gain, the number of antihypertensive medications and LVH.²⁹⁵ Maduell and Navarro (in a cross-sectional study of 15 people) reported that salt restriction alone resulted in a significant reduction in interdialytic weight gain and blood pressure.²⁹⁶ In the Haemodialysis (HEMO) Study, dietary sodium intake was associated with a greater adjusted risk of all-cause mortality.²⁹⁷ In practical terms, adherence may be more sustainable if a threshold restriction of <6 g dietary salt is applied in the diabetes cohort who have additional restrictions placed on them, but this has yet to be formally evaluated.

Diuretic therapy may provide an additional means by which to promote natriuresis in people who are on dialysis who have residual urine output. In the DOPPS study, Bragg-Gresham *et al* reported that diuretic use was associated with reduced interdialytic weight gain, fewer intradialytic hypotensive episodes and reduced cardiac-specific mortality, but not all-cause mortality.²⁸³ People with residual renal function who were on diuretics were twice as likely to retain residual renal function compared with those who were not on diuretics after 1 year in the study. The authors concluded that people with residual renal function may benefit from continuing diuretic use rather than automatically discontinuing it at the start of dialysis. Furthermore, in a prospective randomised study by Medcalf *et al* on people who are on peritoneal dialysis, frusemide given at a dose of 250 mg once daily produced clinically significant preservation in urine volume over 1 year, but it had no influence on residual renal function.²⁹⁸

Interestingly, and independent of its diuretic property, spironolactone has been shown, in a randomised controlled trial that included participants with diabetes, to be more effective than placebo in treating refractory hypertension in people on dialysis.²⁹⁹ There is emerging evidence from a number of randomised controlled trials that spironolactone has a cardiac protective effect in people on dialysis³⁰⁰, but it will be interesting to see what emanates from the current ongoing larger randomised controlled trial (ALdosterone Antagonist Chronic HEModialysis Interventional Survival Trial (ALCHEMIST)), which is exploring the potential cardiac protective role of spironolactone in patients who are on dialysis.³⁰¹ It might help to show whether this effect is dependent or independent of spironolactone's property as a diuretic and/or antihypertensive agent.

To date there have been, therefore, no randomised controlled trials to address the question of whether salt restriction or diuretic use in people with diabetes who are on dialysis may influence blood pressure control or cardiovascular outcome. In the absence of such evidence, individualisation of dietary sodium intake is required, depending on the person's interdialytic weight gain, extracellular volume status, haemodynamic stability and nutritional status.

Declarations of interest

The authors declare the following potential conflicts of interests and support from industry.

- Debasish Banerjee was previously funded for research by the British Heart Foundation (BHF), Kidney Research UK, Wellcome ISSF and received honoraria from AstraZeneca, Pfizer and Vifor Pharma.
- Andrew Frankel has received research grants, and he prepares educational materials and attends drug advisory boards for: Boehringer Ingelheim/Lilly Alliance, AstraZeneca, Novo Nordisk, Merck and Johnson & Johnson.
- Parijat De has received honoraria for educational meetings from Astra Zeneca, Janssen, Boehringer Ingelheim, Novo, Sanofi, Novartis, Abbott, MSD, Takeda, Roche, Lilly, Ascensia, BD, Internis, GSK, Menarini, Bayer and Besins.
- Peter Winocour has received honoraria for delivering educational meetings and/or attending advisory boards for Astra Zeneca, Eli Lilly, Novo Nordisk, Sanofi, MSD, Janssen and Vifor Pharma
- Indranil Dasgupta has received research funding from Medtronic and Sanofi Genzyme, and attended advisory board for GSK

References

- 1 Ritz E, Orth SR. Nephropathy in patients with type 2 diabetes mellitus. *N Engl J Med* 1999;341:1127–33.
- 2 Gilg J, Caskey F, Fogarty D. UK Renal Registry 18th Annual Report: Chapter 1 UK Renal Replacement Therapy Incidence in 2014: National and Centre-specific Analyses. *Nephron* 2016;132(Suppl 1):9–40.
- 3 Stamler J, Vaccaro O, Neaton JD, Wentworth D. Diabetes, other risk factors, and 12-yr cardiovascular mortality for men screened in the Multiple Risk Factor Intervention Trial. *Diabetes Care* 1993;16:434–44.
- 4 Stephens JW, Brown KE, Min T. Chronic kidney disease in type 2 diabetes: Implications for managing glycaemic control, cardiovascular and renal risk. *Diabetes Obes Metab*. 2020 Apr;22 Suppl 1:32–45. doi: 10.1111/dom.13942. PMID: 32267078.
- 5 Rosolowsky ET, Skupien J, Smiles AM *et al*. Risk for ESRD in type 1 diabetes remains high despite renoprotection. *J Am Soc Nephrol* 2011;22:545–53.
- 6 Deckert T, Yokoyama H, Mathiesen E *et al*. Cohort study of predictive value of urinary albumin excretion for atherosclerotic vascular disease in patients with insulin dependent diabetes. *BMJ* 1996;312:871–4.
- 7 Watkins PJ, Parsons V, Bewick M. The prognosis and management of diabetic nephropathy. *Clin Nephrol* 1977;7:243–9.
- 8 Finne P, Reunanen A, Stenman S *et al*. Incidence of end-stage renal disease in patients with type 1 diabetes. *JAMA* 2005;294:1782–7.
- 9 de Boer IH, Rue TC, Cleary PA *et al*. Long-term renal outcomes of patients with type 1 diabetes mellitus and microalbuminuria: an analysis of the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications cohort. *Arch Intern Med* 2011;171:412–420.
- 10 Hollenberg NK, Price DA, Fisher ND *et al*. Glomerular hemodynamics and the renin-angiotensin system in patients with type 1 diabetes mellitus. *Kidney Int* 2003;63:172–8.
- 11 Miller JA. Impact of hyperglycaemia on the renin-angiotensin system in early human type 1 diabetes. *J Am Soc Nephrol* 1999;10:1775–8.
- 12 Lafayette RA, Mayer G, Park SK, Meyer TW. Angiotensin II receptor blockade limits glomerular injury in rats with reduced renal mass. *J Clin Invest* 1992;90:766–71.
- 13 Rosenberg ME, Smith LJ, Correa-Rotter R, Hostetter TH. The paradox of the renin-angiotensin system in chronic renal disease. *Kidney Int* 1994;45:403–10.
- 14 Ray PE, Aguilera G, Kopp JB *et al*. Angiotensin II receptor-mediated proliferation of cultured human fetal mesangial cells. *Kidney Int* 1991;40:764–71.
- 15 Harindhanavudhi T, Mauer M, Klein R *et al*. Benefits of Renin-Angiotensin blockade on retinopathy in type 1 diabetes vary with glycemic control. *Diabetes Care* 2011;34:1838–42.
- 16 Sochett EB, Cherney DZ, Curtis JR *et al*. Impact of renin angiotensin system modulation on the hyperfiltration state in type 1 diabetes. *J Am Soc Nephrol* 2006;17:1703–9.
- 17 Raile K, Galler A, Hofer S *et al*. Diabetic nephropathy in 27,805 children, adolescents, and adults with type 1 diabetes: effect of diabetes duration, A1C, hypertension, dyslipidemia, diabetes onset, and sex. *Diabetes Care* 2007;30:2523–8.
- 18 Chowdhury TA, Kumar S, Barnett AH, Bain SC. Nephropathy in type 1 diabetes: the role of genetic factors. *Diabet Med* 1995;12:1059–68.

- 19 Microalbuminuria Collaborative Study Group, United Kingdom. Risk factors for development of microalbuminuria in insulin dependent diabetic patients: a cohort study. *BMJ* 1993;306:1235–9.
- 20 Rudberg S, Dahlquist G. Determinants of progression of microalbuminuria in adolescents with IDDM. *Diabetes Care* 1996;19:369–71.
- 21 Ku E, McCulloch CE, Mauer M *et al.* Association between blood pressure and adverse renal events in type 1 diabetes. *Diabetes Care* 2016;30:2218–24.
- 22 Guo J, Brooks M, Muldoon M *et al.* Optimal blood pressure thresholds for minimal coronary artery disease risk in type 1 diabetes. *Diabetes Care* 2019; 2:1692–9.
- 23 Kidney Disease: Improving Global Outcomes (KDIGO) Clinical practice guideline on the management of blood pressure in chronic kidney disease. Public review draft January 2020. <https://kdigo.org/wp-content/uploads/2016/10/KDIGO-BP-Management-in-CKD-GL-public-review-draft-Jan-31-2020.pdf>
- 24 National Institute for Health and Care Excellence. *Type 1 diabetes in adults: diagnosis and management*, NICE guideline NG17, 2016. Available from: www.nice.org.uk/guidance/ng17 [Accessed February 2017].
- 25 Tuttle KR, Bakris GL, Bilous RW *et al.* Diabetic kidney disease: a report from an ADA Consensus Conference. *Am J Kidney Dis* 2014;64:510–33.
- 26 Donaghue KC, Wadwa RP, Dimeglio LA *et al.* ISPAD Clinical Practice Consensus Guidelines 2014. Microvascular and macrovascular complications in children and adolescents. *Pediatr Diabetes* 2014;15(Suppl 20):257–69.
- 27 Guo J, Muldoon M, Brooks M, Orchard T, Costacou T. Prognostic significance of pulse pressure and other blood pressure components for coronary artery disease in type 1 diabetes. *American Journal of Hypertension* 2009;32:1075–81.
- 28 Sinclair AJ, Dhatriya K. Clinical guidelines for type 1 diabetes mellitus with an emphasis on older adults: an executive summary. *Diabet Med* 2020; 37:53–70.
- 29 2019 ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD: The Task Force for diabetes, pre-diabetes, and cardiovascular diseases of the European Society of Cardiology (ESC) and the European Association for the Study of Diabetes (EASD). *European Heart Journal* 2020;41:255–323.
- 30 Lurbe E, Redon J, Kesani A *et al.* Increase in nocturnal blood pressure and progression to microalbuminuria in type 1 diabetes. *N Engl J Med* 2002;347:797–805.
- 31 Shalaby NM, Shalaby NM. Study of ambulatory bloods pressure in diabetic children: prediction of early renal insult. *Ther Clin Risk Manag* 2015;11:1531–7.
- 32 Lambers Heerspink HJ, Holtkamp FA, Parvin HH *et al.* Moderation of dietary sodium potentiates the renal and cardiovascular protective effects of angiotensin receptor blockers. *Kidney Int* 2012;82:330–7.
- 33 Lithovius R, Harjutsalo V, Forsblom C *et al.* Antihypertensive treatment and resistant hypertension in patients with type 1 diabetes by stages of diabetic nephropathy. *Diabetes Care* 2014;37:709–17.
- 34 National Institute for Health and Care Excellence. *Hypertension in adults: diagnosis and management*, NICE guideline NG136, 2019. Available from: www.nice.org.uk/guidance/CG127 [Accessed September 2020].
- 35 Alderman MH. Salt, Blood Pressure and Human Health. *Hypertension* 2000;36:890–3.
- 36 Jones DW, Kim JS, Andrew ME *et al.* Body mass index and blood pressure in Korean men and Women: the Korean National Blood Pressure Survey. *J Hypertens* 1994;12:1433–7.

- 37 Whelton SP, Chin A, Xin X, HE J. Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. *Ann Intern Med* 2002;136:493–503.
- 38 Chen L, Smith GD, Harbord RM, Lewis SJ. Alcohol intake and blood pressure: a systematic review implementing a Mendelian randomization approach. *PLoS Med* 2008;5:e52.
- 39 Gerds E, Svarstad E, Myking OL, Lund-Johansen P, Omvik P. Salt sensitivity in hypertensive type-1 diabetes mellitus. *Blood Press* 1996;5:78–85.
- 40 Thomas MC, Moran J, Forsblom C *et al.* The association between dietary sodium intake, ESRD, and all-cause mortality in patients with type 1 diabetes. *Diabetes Care* 2011;34:861–866.
- 41 Dickinson HO, Mason JM, Nicolson DJ *et al.* Lifestyle interventions to reduce raised blood pressure: a systematic review of randomized controlled trials. *J Hypertens* 2006;24:215–33.
- 42 Slagman MC, Waanders F, Hemmelder MH *et al.* Moderate dietary sodium restriction added to angiotensin converting enzyme inhibition compared with dual blockade in lowering proteinuria and blood pressure: randomised controlled trial. *BMJ* 2011;343:d4366.
- 43 Vogt L, Waanders F, Boomsma F *et al.* Effects of Dietary Sodium and Hydrochlorothiazide on the Antiproteinuric Efficacy of Losartan. *J Am Soc Nephrol* 2008;19:999–1007.
- 44 Melin EO, Thulesius HO, Hillman M, Landin-Olsson M and Thunander M. Abdominal obesity in type 1 diabetes associated with gender, cardiovascular risk factors and complications, and difficulties achieving treatment targets: a cross sectional study at a secondary care diabetes clinic *BMC Obesity* 2018;5:15.
- 45 Purnell JQ, Hokanson JE, Marcovina SM *et al.* Effect of excessive weight gain with intensive therapy of type 1 diabetes on lipid levels and blood pressure results from the DCCT. *JAMA* 1998;280:140–6.
- 46 Navaneethan SD, Yehnert H, Moustarah F *et al.* Weight loss interventions in chronic kidney disease. A systematic review and meta-analysis. *Clin J Am Soc Nephrol* 2009;4:1565–74.
- 47 Ostman C, Jewiss D, King N, Smart NA. Clinical outcomes to exercise training in type 1 diabetes: A systematic review and meta-analysis. *Diabetes Res Clin Pract* 2018;139:380–91.
- 48 Chen JL, Lerner D, Ruthazer R *et al.* Association of physical activity with mortality in chronic kidney disease. *J Nephrol* 2008;21:243–52.
- 49 World Health Organization. *International guide for monitoring alcohol consumption and related harm*. Geneva: WHO, 2000.
- 50 The EUCLID Study Group. Randomised placebo-controlled trial of lisinopril in normotensive patients with insulin dependent diabetes and normoalbuminuria or microalbuminuria. *Lancet* 1997;349:1787–92.
- 51 Bilous R, Chaturvedi N, Sjølie AK *et al.* Effect of candesartan on microalbuminuria and albumin excretion rate in diabetes: three randomized trials. *Ann Intern Med* 2009;151:11–20.
- 52 Mauer M, Zinman B, Gardiner R *et al.* Renal and retinal effects of enalapril and losartan in type 1 diabetes. *N Engl J Med* 2009;361:40–51.
- 53 Kventy J, Gregersen G, Smith Pedersen R. Randomized placebo-controlled trial of perindopril in normotensive, normoalbuminuric patients with type 1 diabetes mellitus. *QJM* 2001;94:89–94.

- 54 Almdal T, Norgaard K, Feldt-Rasmussen B, Deckert T. The predictive value of microalbuminuria in IDDM. A five-year follow-up study. *Diabetes Care* 1994;17:120–5.
- 55 Hovind P, Tarnow L, Rossing P *et al.* Predictors for the development of microalbuminuria and macroalbuminuria in patients with type 1 diabetes: inception cohort study. *BMJ* 2004;328:1105–9.
- 56 Perkins BA, Ficociello LH, Silva KH *et al.* Regression of microalbuminuria in type 1 diabetes. *N Engl J Med* 2003;348:2285–93.
- 57 Forsblom CM, Groop PH, Ekstrand A, Groop LC. Predictive value of microalbuminuria in patients with insulin-dependent diabetes of long duration. *BMJ* 1992;305:1051–3.
- 58 de Boer IH, Afkarian M, Rue TC *et al.* Renal outcomes in patients with type 1 diabetes and macroalbuminuria. *J Am Soc Nephrol* 2014;25:2342–50.
- 59 Macisaac RJ, Jerums G. Diabetic kidney disease with and without albuminuria. *Curr Opin Nephrol Hypertens* 2011;20:246–57.
- 60 Perkins BA, Ficociello LH, Roshan B *et al.* In patients with type 1 diabetes and new-onset microalbuminuria the development of advanced chronic kidney disease may not require progression to proteinuria. *Kidney Int* 2010;77:57–64.
- 61 Dunger DB, Schwarze CP, Cooper JD *et al.* Can we identify adolescents at high risk for nephropathy before the development of microalbuminuria? *Diabet Med* 2007;24:131–6.
- 62 Maftai O, Pena AS, Sullivan T *et al.* Early atherosclerosis relates to urinary albumin excretion and cardiovascular risk factors in adolescents with type 1 diabetes: Adolescent type 1 Diabetes cardio-renal Intervention Trial (AddIT). *Diabetes Care* 2014;37:3069–75.
- 63 Krolewski AS. Progressive Renal Decline: The New Paradigm of Diabetic Nephropathy in Type 1 Diabetes. *Diabetes Care* 2015;38:954–62.
- 64 Rahimi Z, Moradi M, Nasri H. A systematic review of the role of renin angiotensin aldosterone system genes in diabetes mellitus, diabetic retinopathy and diabetic neuropathy. *J Res Med Sci* 2014;19:1090–8.
- 65 Krolewski AS, Warram JH, Forsblom C *et al.* Serum concentration of cystatin C and risk of end-stage renal disease in diabetes. *Diabetes Care* 2012;35:2311–6.
- 66 Gohda T, Niewczas MA, Ficociello LH *et al.* Circulating TNF receptors 1 and 2 predict stage 3 of CKD in type 1 diabetes. *J Am Soc Nephrol* 2012;23:516–524.
- 67 Cantarovich D, Perrone V. Pancreas transplant as treatment to arrest renal function decline in patients with type 1 diabetes and proteinuria. *Semin Nephrol* 2012;32:432–6.
- 68 Mathiesen ER, Hommel E, Hansen HP *et al.* Randomised controlled trial of long term efficacy of captopril on preservation of kidney function in normotensive patients with insulin dependent diabetes and microalbuminuria. *BMJ* 1999;319:24–5.
- 69 The Microalbuminuria Collaborative Study Group. Captopril reduces the risk of nephropathy in IDDM patients with microalbuminuria. *Diabetologia* 1996;35:587–93.
- 70 O'Hare P, Bilbous R, Mitchell T *et al.* Low-dose ramipril reduces microalbuminuria in type 1 diabetic patients without hypertension: results of a randomized controlled trial. *Diabetes Care* 2000;23:1823–9.
- 71 Marre M, Chatellier G, Leblanc H *et al.* Prevention of diabetic nephropathy with enalapril in normotensive diabetics with microalbuminuria. *BMJ* 1998;297:1092–5.
- 72 Viberti G, Mogensen CE, Groop LC, Pauls JF. Effect of captopril on progression to clinical proteinuria in patients with insulin-dependent diabetes mellitus and

- microalbuminuria. European Microalbuminuria Captopril Study Group. *JAMA* 1994;271:275–279.
- 73 The ACE Inhibitors in Diabetic Nephropathy Trialist Group. Should all patients with type 1 diabetes mellitus and microalbuminuria receive angiotensin-converting enzyme inhibitors? A meta-analysis of individual patient data. *Ann Intern Med* 2001;134:370–9.
 - 74 Parving HH, Andersen AR, Smidt UM, Svendsen PA. Early aggressive antihypertensive treatment reduces rate of decline in kidney function in diabetic nephropathy. *Lancet* 1983;8335:1175–9.
 - 75 Parving HH, Hommel, Smidt U. Protection of kidney function and decrease in albuminuria by captopril in insulin dependent diabetics with nephropathy. *BMJ* 1988;297:1086–91.
 - 76 Lewis EJ, Hunsicker LG, Bain RP, Rohde RD. The effect of angiotensin-converting enzyme inhibition on diabetic nephropathy. *N Eng J Med* 1993;1456–62.
 - 77 Björck S, Mulec H, Johnsen SA *et al*. Renal protective effect of enalapril in diabetic nephropathy. *BMJ* 1992;304:339–43.
 - 78 Sarafidis PA, Stafylas PC, Kanaki AI, Lasaridis AN. Effects of renin-angiotensin system blockers on renal outcomes and all-cause mortality in patients with diabetic nephropathy: an updated meta-analysis. *Am J Hypertens* 2008;21:922–9.
 - 79 Chaturvedi N, Porta M, Klein R *et al*. Effect of candesartan on prevention (DIRECT-Prevent 1) and progression (DIRECT-Protect 1) of retinopathy in type 1 diabetes: randomised, placebo-controlled trials. *Lancet* 2008;372:1394–402.
 - 80 Andersen S, Tarnow L, Rossing P *et al*. Renoprotective effects of angiotensin II receptor blockade in type 1 diabetic patients with diabetic nephropathy. *Kidney Int* 2000;57:601–6.
 - 81 Schjoedt KJ, Andersen S, Rossing P *et al*. Aldosterone escape during blockade of the renin-angiotensin-aldosterone system in diabetic nephropathy is associated with enhanced decline in glomerular filtration rate. *Diabetologia* 2004;47:1936–9.
 - 82 Nielsen SE, Persson F, Frandsen E *et al*. Spironolactone diminishes urinary albumin excretion in patients with type 1 diabetes and microalbuminuria: a randomized placebo-controlled crossover study. *Diabetic Med* 2012;29:e184–90.
 - 83 Heerspink HJ, Persson F, Brenner BM *et al*. Renal outcomes with aliskiren in patients with type 2 diabetes: a prespecified secondary analysis of the ALTITUDE randomised controlled trial. *Lancet Diabetes Endocrinol* 2016;4:309–17.
 - 84 Cherney DZ, Scholey JW, Jiang S *et al*. The effect of direct renin inhibition alone and in combination with ACE inhibition on endothelial function, arterial stiffness, and renal function in type 1 diabetes. *Diabetes Care* 2012;35:2324–30.
 - 85 Cherney DZ, Lai V, Scholey JW *et al*. Effect of direct renin inhibition on renal hemodynamic function, arterial stiffness, and endothelial function in humans with uncomplicated type 1 diabetes: a pilot study. *Diabetes Care* 2010;33:361–5.
 - 86 Elving LD, Wetzels JFM, van Lier HJJ *et al*. Captopril and atenolol are equally effective in retarding progression of diabetic nephropathy. Results of a 2 year randomised study. *Diabetologia* 1994;37:604–9.
 - 87 Bakris GL, Weir MR, Secic M *et al*. Differential effects of calcium antagonist subclasses on markers of nephropathy progression. *Kidney Int* 2004;65:1991–2002.
 - 88 Cherney DZ, Perkins BA, Soleymanlou N *et al*. Renal hemodynamic effect of sodium-glucose cotransporter 2 inhibition in patients with type 1 diabetes mellitus. *Circulation* 2014;129:587–97.

- 89 Wanner C, Inzucchi SE, Lachin JM *et al.* Empagliflozin and Progression of Kidney Disease in Type 2 Diabetes. *N Engl J Med* 2016;375:323–34.
- 90 Van Raalte DH, Bjornstad P, Persson F *et al.* The impact of Sotagliflozin on renal function, albuminuria, and blood pressure in adults with type 1 diabetes (T1D). *Diabetes* 2019;68(Supplement 1)
- 91 Peters AL, Buschur EO, Buse JB *et al.* Euglycemic Diabetic Ketoacidosis: A Potential Complication of Treatment With Sodium-Glucose Cotransporter 2 Inhibition. *Diabetes Care* 2015;38:1687–93.
- 92 Jennings DL, Kalus JS, Coleman CI *et al.* Combination therapy with an ACE inhibitor and an angiotensin receptor blocker for diabetic nephropathy: a meta-analysis. *Diabet Med* 2007;24:486–93.
- 93 Pham JT, Schmitt BP, Leehey DJ. Effects of dual blockade of the renin-angiotensin system in diabetic kidney disease: a systematic review and meta-analysis. *J Nephrol Therapeut* 2012;(Suppl 2):003.
- 94 Maione A, Navaneethan SD, Graziano G *et al.* Angiotensin-converting enzyme inhibitors, angiotensin receptor blockers and combined therapy in patients with micro- and macroalbuminuria and other cardiovascular risk factors: a systematic review of randomized controlled trials. *Nephrol Dial Transplant* 2011;26:2827-47.
- 95 Fried LF, Emanuele N, Zhang JH *et al.* Combined angiotensin inhibition for the treatment of diabetic nephropathy. *N Engl J Med* 2013;369:1892–903.
- 96 Anantharaman R, Bhansali A, Bhadada SK *et al.* Anti-albuminuric efficacy of a combination of angiotensin converting enzyme inhibitor & angiotensin receptor blocker in type 1 DM with nephropathy. *Indian J Med Res* 2010;132:42–7.
- 97 Jacobsen P, Andersen S, Rossing K *et al.* Dual blockade of the renin-angiotensin system in type 1 patients with diabetic nephropathy. *Nephrol Dial Transplant* 2002;17:1019–24.
- 98 Cooper WO, Hernandez-Diaz S, Arbogast PG *et al.* Major congenital malformations after first-trimester exposure to ACE inhibitors. *N Engl J Med* 2006;354:2443–51.
- 99 Li D, Yang C, Andrade S *et al.* Maternal exposure to angiotensin converting enzyme inhibitors in the first trimester and risk of malformations in offspring: a retrospective cohort study. *BMJ* 2011;343:d5931.
- 100 Bakris GL, Weir MR. Angiotensin-converting enzyme inhibitor-associated elevations in serum creatinine: is this a cause for concern? *Arch Intern Med* 2000;160:685–93.
- 101 Postma CT, Klappe EM, Dekker HM, Thien T. The prevalence of renal artery stenosis among patients with diabetes mellitus. *Eur J Intern Med* 2012;23:639–42.
- 102 Chowdhury TA. Taking precautions with angiotensin converting enzyme inhibitors. Clinically significant deterioration in renal function occurs rarely (letter). *BMJ* 1999;318:258.
- 103 Reams GP, Bauer JH, Gaddy P. Use of the converting enzyme inhibitor enalapril in renovascular hypertension. Effect on blood pressure, renal function, and the renin-angiotensin-aldosterone system. *Hypertension* 1986;8:290–7.
- 104 Palmer BF. Managing hyperkalemia caused by inhibitors of the renin-angiotensin-aldosterone system. *N Engl J Med* 2004;351:585–92.
- 105 Anonymous. Sick day rules in kidney disease. *Drug and Therapeutics Bulletin* 2015;53:37.
- 106 Rossing P. Diabetic Nephropathy: worldwide epidemic and effects of current treatment on natural history. *Curr Diab Rep* 2006;6:479–483.

- 107 Schena FP, Gesualdo L. Pathogenic mechanisms of diabetic nephropathy. *J Am Soc Nephrol* 2005;16(Suppl 1):S30–3.
- 108 Adler S. Diabetic nephropathy: Linking histology, cell biology, and genetics. *Kidney Int* 2004;66:2095–106.
- 109 Ritz E, Rychlik I, Locatelli F, Halimi S. End-stage renal failure in type 2 diabetes: A medical catastrophe of worldwide dimensions. *Am J Kidney Dis* 1999;34:795–808.
- 110 Adler AI, Stevens RJ, Manley SE *et al*. Development and progression of nephropathy in type 2 diabetes: the United Kingdom Perspective Diabetes Study (UKPDS 64). *Kidney Int* 2003;63:225–32.
- 111 Angélique ME, Spoelstra-de Man, Catherine B *et al*. Rapid Progression of Albumin Excretion is an Independent Predictor of Cardiovascular Mortality in Patients with Type 2 Diabetes and Microalbuminuria. *Diabetes Care* 2001;24:2097–101.
- 112 Ritz E, Dikow R. Hypertension and antihypertensive treatment of diabetic nephrology. *Nat Clin Pract Nephrol* 2006;2:562–567.
- 113 Parving HH, Osterby R, Ritz E. Diabetic Nephropathy. In: Brenner BM, Levine S, editors. *The Kidney*. 6th ed. Philadelphia: WB Saunders, 2000: 1731–73.
- 114 Klag MJ, Whelton PK, Randall BL *et al*. Blood pressure and end-stage renal disease in men. *N Engl J Med* 1996;334:13–18.
- 115 O'Seaghdha CM, Perkovic V, Lam TH *et al*. Blood pressure is a major risk factor for renal death: an analysis of 560 352 participants from the Asia-Pacific region. *Hypertension* 2009;54:509–15.
- 116 Singh R, Singh AK, Alavi N, Leehey DJ. Mechanism of increased angiotensin II levels in glomerular mesangial cells cultured in high glucose. *J Am Soc Nephrol* 2003;14:873–80.
- 117 American Diabetes Association: Hypertension Management in Adults with Diabetes (position statement). *Diabetes Care* 2004;27(Suppl 1):S65–7.
- 118 Tuttle KR, Bakris GL, Bilous RW *et al*. Diabetic kidney disease: a report from an ADA Consensus Conference. *Diabetes Care* 2014;37:2864–83.
- 119 Beckett NS, Peters R, Fletcher AE *et al*. Treatment of hypertension in patients 80 years of age or older. *N Engl J Med* 2008;358:1887–98.
- 120 Lindholm LH, Hansson L, Ekblom T *et al* for the STOP Hypertension-2 Study Group. Comparison of antihypertensive treatments in preventing cardiovascular events in elderly diabetic patients: results from the Swedish Trial in Old Patients with Hypertension –2. *J Hypertens* 2000;18:1671–75.
- 121 Agarwal R, Andersen MJ. Prognostic Importance of clinic and home blood pressure recordings in patients with chronic kidney disease. *Kidney Int* 2006;69:406–11.
- 122 Agarwal R, Andersen MJ. Blood pressure recordings within and outside the clinic and cardiovascular events in chronic kidney disease. *Am J of Nephrol* 2006;26:503–10.
- 123 Davidson MB, Hix JK, Vidt DG, Brotman DJ. Association of impaired diurnal blood pressure variation with subsequent decline in glomerular filtration rate. *Arch Intern Med* 2006;166:846–52.
- 124 Minutolo R, Agarwal R, Borrelli S *et al*. Prognostic role of ambulatory blood pressure measurement in patients with non-dialysis kidney disease. *Arch Intern Med* 2011;171:1090–8.
- 125 Ravid, M, Lang R, Rachmani R, Lishner M. Long-term renoprotective effect of angiotensin-converting enzyme inhibition in non-insulin dependent diabetes mellitus: A 7-year follow-up study. *Arch Intern Med* 1996;156:286–9.

- 126 Parving HH, Lehnert H, Bröchner-Mortensen J *et al.* The effect of irbesartan on the development of diabetic nephropathy in patients with type 2 diabetes. *N Eng J Med* 2001;345:870–8.
- 127 Brenner BM, Cooper ME, de Zeeuw D *et al.* Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001;345:861–9.
- 128 Rossing P, Hommel E, Schmidt UM, Parving HH. Reduction in albuminuria predicts a beneficial effect on diminishing the progression of human diabetic nephropathy during antihypertensive treatment. *Diabetologia* 1994;37:511–6.
- 129 Lawson A, Hameed MA, Brown R *et al.* Non-adherence to antihypertensive medications is related to pill burden in apparent treatment resistant hypertensive individuals. *J Hypertens* 2020;38:1165–73.
- 130 Heart Outcomes Prevention Evaluation Study Investigators. Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: results of the HOPE study and the MICRO-HOPE substudy. *Lancet* 2000;355:253–9.
- 131 Standards of Medical Care in Diabetes – 2015: Summary of Revisions. *Diabetes Care* 2015;38(Suppl 1):S4.
- 132 Weber MA, Schiffrin EL, White WB *et al.* Clinical practice guidelines for the management of hypertension in the community. A statement by the American Society of Hypertension and the International Society of Hypertension. *J Hyperten* 2014;32:3–15.
- 133 KDOQI. Clinical Practice Guidelines and Clinical Practice Recommendations for Diabetes and Chronic Kidney Disease. *Am J Kid Dis* 2007;49(Suppl 2):S12–154.
- 134 Mancia G, Fagard R, Narkiewicz K *et al.* 2013 ESH/ESC Guidelines for the management of arterial hypertension: the Task Force for the management of arterial hypertension of the European Society of Hypertension (ESH) and the European Society of Cardiology (ESC). *J Hypertens* 2013;31:1281–357.
- 135 James PA, Oparil S, Carter BL *et al.* 2014 evidence-based guidance for the management of high blood pressure in adults: report from the panel members appointed to the eighth Joint National Committee (JNC 8). *JAMA* 2014;311:507–20.
- 136 Bangalore S, Fakheri R, Toklu B, Messerli FH. Diabetes mellitus as a compelling indication for use of renin angiotensin system blockers: systematic review and meta-analysis of randomized trials. *BMJ* 2016;352:i438.
- 137 National Institute for Health and Care Excellence. *Chronic kidney disease in adults: assessment and management*, NICE guideline CG182, issued July 2014 and modified March 2015.
- 138 Casas JP, Chua W, Loukogeorgakis S *et al.* Effects of inhibitors of the renin-angiotensin system and other antihypertensive drugs on renal outcomes: system review and meta-analysis. *Lancet* 2005;366:2026–33.
- 139 Lewis EJ, Hunsicker LG, Clarke WR *et al.* Renoprotective effect of angiotensin-receptor antagonist irbesartan in patients with nephropathy due to type 2 diabetes. *N Eng J Med* 2001;345:851–60.
- 140 Barnett AH, Bain SC, Bouter P *et al.* Angiotensin-receptor blocker versus converting-enzyme inhibition in type 2 diabetes and nephropathy. *N Eng J Med* 2004;351:1952–61.
- 141 Ennezat PV, Berlowitz M, Sonnenblick EH, Le Jemtel DH. Therapeutic implications of escape from angiotensin-converting enzyme inhibition in patients with chronic heart failure. *Curr Cardiol Rep* 2000;2:258–62.

- 142 Balcells E, Meng QC, Johnson WH Jr *et al.* Angiotensin II formation from ACE and chymase in human and animal hearts: methods and species considerations. *Am J Physiol* 1997;273:H1769–74.
- 143 Mogensen CE, Neldam S, Tikkanen I *et al.* Randomised controlled trial of dual blockade of renin-angiotensin system in patients with hypertension, microalbuminuria, and non-insulin dependent diabetes: the candesartan and lisinopril microalbuminuria (CALM) study. *BMJ* 2000;321:1440–4.
- 144 Mann JF, Schmieder RE, McQueen M *et al.* Renal outcomes with telmisartan, ramipril, or both, in people at high vascular risk (the ONTARGET study): a multicentre, randomised, double-blind, controlled trial). *Lancet* 2008;372:547–53.
- 145 Perkovic V, Agarwal R, Fioretto P *et al.* Management of patients with diabetes and CKD: conclusions from a ‘Kidney Disease: Improving Global Outcomes’ (KDIGO) Controversies Conference. *Kidney Int* 2016;90:1175–83
- 146 Biesen WV, Fouque D, Wiecek A, Cochat P *et al.* A European renal best practice position statement on kidney disease and endorsement of KDIGO guidelines with some caveats for real life application. *Nephrol Dial Transplant* 2013;0:1–8.
- 147 Campbell NR, Khan NA, Hill MD *et al.* 2009 Canadian Hypertension Education Program Recommendations: the scientific summary – an annual update. *Can J Cardiol* 2009;25:271–7.
- 148 Dalla Vestra M, Simioni N, Masiero A. Renal effects of dual renin-angiotensin-aldosterone system blockade in patients with diabetic nephropathy. *Int Urol Nephrol* 2009;41:119–126.
- 149 Phillips CO, Kashani A, Ko DK, Francis G, Krumholz HM. Adverse effects of combination angiotensin II receptor blockers plus angiotensin-converting enzyme inhibitors for left ventricular dysfunction: a quantitative review of data from randomized clinical trials. *Arch Intern Med* 2007;167:1930–6.
- 150 Rocha R, Stier CT Jr, Kifor I *et al.* Aldosterone: a mediator of myocardial necrosis and renal arteriopathy. *Endocrinology* 2000;141:3871–8.
- 151 Struthers AD. Aldosterone: cardiovascular assault. *Am Heart J* 2002;144:S2–7.
- 152 Sato A, Hayashi K, Naruse M, Saruta T. Effectiveness of aldosterone blockade in patients with diabetic nephropathy. *Hypertension* 2003;41:64–8.
- 153 Rachmani R, Slavachevsky I, Amit M *et al.* The effect of spironolactone, cilazapril and their combination on albuminuria in patients with hypertension and diabetic nephropathy is independent of blood pressure reduction: a randomized controlled study. *Diabet Med* 2004;21:471–5.
- 154 Epstein M, Buckalew V, Altamirano J *et al.* Eplerenone reduces proteinuria in type II diabetes: Implications for aldosterone involvement in the pathogenesis of renal dysfunction (Abstract). *J Am Coll Cardiol* 2002;39(Suppl A):249A.
- 155 Epstein M, Williams GH, Weinberger M *et al.* Selective aldosterone blockade with eplerenone reduces albuminuria in patients with type 2 diabetes. *Clin J Am Soc Nephrol* 2006;1:940–51.
- 156 Bakris GL, Agarwal R, Chan JC, Cooper ME *et al.* Effect of finerenone on albuminuria in patients with diabetic nephropathy: a randomized clinical trial. *JAMA* 2015;314:884–94.
- 157 Bakris GL, Agarwal R, Anker SD *et al.* on behalf of the FIDELIO-DKD study investigators. Design and Baseline Characteristics of the Finerenone in Reducing Kidney Failure and Disease Progression in Diabetic Kidney Disease Trial. *Am J Nephrol* 2019;50:333–344.

- 158 Bakris GL, Agarwal R, Anker SD *et al* for the FIDELIO-DKD study investigators. Effect of finerenone on chronic kidney disease outcomes in type 2 diabetes. *New Eng J Med* October 23, 2020. DOI: 10.1056/NEJMoa2025845
- 159 Parving HH, Brenner BM, McMurrige AJ *et al*. Aliskiren trial in type 2 diabetes using cardiorenal end-points (ALTITUDE): Rationale and study design. *Nephrol Dial Transplant* 2009;24:1663–71.
- 160 Parving HH, Persson F, Lewis JB *et al*. Aliskiren combined with losartan in type 2 diabetes and nephropathy. *N Eng J Med* 2008;358:2433–46.
- 161 National Institute for Health and Care Excellence. *Patiromer for treating hyperkalaemia*. Technology appraisal guidance [TA623]. NICE, 2020. Available from: www.nice.org.uk/guidance/ta623/chapter/1-Recommendations [Accessed: 6 June 2020].
- 162 National Institute for Health and Care Excellence. *Sodium zirconium cyclosilicate for treating hyperkalaemia*. Technology appraisal guidance [TA599]. NICE, 2019. Available from: www.nice.org.uk/guidance/ta599/chapter/1-Recommendations [Accessed: 6 June 2020].
- 163 Kidney Disease Outcomes Quality Initiative (KDOQI). KDOQI Clinical Practice Guidelines on hypertension and antihypertensive agents in Chronic Kidney Disease. *Am J Kid Dis* 2004;43(5 Suppl 1):S1–290.
- 164 Weir MR. Acute fall in glomerular filtration rate with renin-angiotensin system inhibition: a biomeasure of therapeutic success? *Kidney Int* 2011;80:235–7.
- 165 US Food and Drug Administration. Public Health Advisory: Angiotensin-converting enzyme inhibitor (ACE inhibitor) drugs and pregnancy, 2006. Available from www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm053113.htm [Accessed February 2017].
- 166 Palmer BF. Renal dysfunction complicating the treatment of hypertension. *N Eng J Med* 2002;347:1256–61.
- 167 Bridoux F, Hazzan M, Pallot JL *et al*. Acute renal failure after the use of angiotensin-converting enzyme inhibitors in patients without renal artery stenosis. *Nephrol Dial Transplant* 1992;7:100–104.
- 168 Kohli HS, Bhaskaran MC, Muthukumar T *et al*. Treatment-related acute renal failure in the elderly: a hospital-based prospective study. *Nephrol Dial Transplant* 2000;15:212–7.
- 169 Smith AC, Toto R, Bakris GL. Differential effects of calcium channel blockers on size selectively of proteinuria in diabetic glomerulopathy. *Kidney Int* 1998;54:889–96.
- 170 Toto RD, Tian M, Fakouhi K *et al*. Effects of calcium channel blockers on proteinuria in patients with diabetic nephropathy. *J Clin Hypertens* 2008;10:761–9.
- 171 Neumann J, Ligtenberg G, Klein II *et al*. Sympathetic hyperactivity in chronic kidney disease: pathogenesis, clinical relevance, and treatment. *Kidney Int* 2004;65:1568–76.
- 172 Bleyer AJ, Hartman J, Brannon PC *et al*. Characteristics of sudden death in haemodialysis patients. *Kidney Int* 2006;69:2268–73.
- 173 Badve SV, Roberts MA, Hawley CM *et al*. Effects of beta-adrenergic antagonists in patients with chronic kidney disease: a systematic review and meta-analysis. *J Am Coll Cardiol* 2011;58:1152–61.
- 174 ALLHAT Officers and Coordinators for the ALLHAT Collaborative Research Group. Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting inhibitor or calcium channel blocker vs diuretic: The Anti-hypertensive and

- Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA* 2002;288:2981–97.
- 175 PROGRESS Collaborative Group. Randomised trial of a perindopril-based blood-pressure-lowering regimen among 6,105 individuals with previous stroke on transient ischaemic attack. *Lancet* 2001;358:1033–41.
 - 176 Vonend O, Marsalek P, Russ H *et al*. Moxonidine treatment of hypertensive patients with advanced renal failure. *J Hypertens* 2003;21:1709–17.
 - 177 Heerspink HJL, Andress DL, Bakris G *et al*. Baseline characteristics and enrichment results from the SONAR trial. *Diabetes Obes Metab* 2018;20:1829–35.
 - 178 Perkovic V, Jardine MJ, Neal B *et al* for the CREDENCE Trial Investigators. Canagliflozin and Renal Outcomes in Type 2 Diabetes and Nephropathy. *N Engl J Med* 2019; 380:2295–2306.
 - 179 Canagliflozin and Cardiovascular and Renal Outcomes in Type 2 Diabetes Mellitus and Chronic Kidney Disease in Primary and Secondary Cardiovascular Prevention Groups. Results from the randomized CREDENCE Trial. *Circulation* 2019;140:739–50.
 - 180 Ingelfinger JR, Rosen CJ. Finerenone. Halting relative hyperaldosteronism in chronic kidney disease. *N Engl J Med* 2020;383:2285–6. doi: 10.1056/NEJMe2031382.
 - 181 Einhorn LM, Zhan M, Hsu VD *et al*. The frequency of hyperkalemia and its significance in chronic kidney disease. *Arch Intern Med* 2009;169:1156–62.
 - 182 Loutradis C, Tolika P, Skodra A *et al*. Prevalence of Hyperkalemia in Diabetic and Non-Diabetic Patients with Chronic Kidney Disease: A Nested Case-Control Study. *Am J Nephrol* 2015;42:351–60.
 - 183 Weinberg JM, Appel LJ, Bakris G *et al*. Risk of hyperkalemia in nondiabetic patients with chronic kidney disease receiving antihypertensive therapy. *Arch Intern Med* 2009;169:1587–94.
 - 184 Umpierrez GE, Cantey P, Smiley D, Palacio A, Temponi D, Luster K, Chapman A. Primary aldosteronism in diabetic subjects with resistant hypertension. *Diabetes Care* 2007;30:1699–1703.
 - 185 Adler AI, Stratton IM, Neil HA *et al*. Association of systolic blood pressure with macrovascular and microvascular complications of type 2 diabetes (UKPDS 36): prospective observational study. *BMJ* 2000;321:412–19.
 - 186 UK Prospective Diabetes Study Group. Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. *BMJ* 1998;317:703–13.
 - 187 Hansson L, Zanchetti A, Carruthers SG *et al*. Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. HOT Study Group. *Lancet* 1998;351:1755–62.
 - 188 Brunström M, Carlberg B. Effect of antihypertensive treatment at different blood pressure levels in patients with diabetes mellitus: systematic review and meta-analyses. *BMJ* 2016;352:i717.
 - 189 Takahashi S, Katada J, Daida H *et al*. Effects of mineralocorticoid receptor antagonists in patients with hypertension and diabetes mellitus: a systematic review and meta-analysis. *J Hum Hypertens* 2016;30:534–42.
 - 190 Lambers Heerspink HJ, Tighiouart H, Sang Y *et al*. GFR decline and subsequent risk of established kidney outcomes: a meta-analysis of 37 randomized controlled trials. *Am J Kidney Dis* 2014;64:860–6.

- 191 ACCORD Study Group, Cushman WC, Evans GW *et al.* Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010;362:1575–85.
- 192 SPRINT Research Group, Wright JT Jr, Williamson JD *et al.* A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N Engl J Med* 2015;373:2103–16.
- 193 Banerjee D, Brincat S, Gregson H *et al.* Pulse pressure and inhibition of renin-angiotensin system in chronic kidney disease. *Nephrol Dial Transplant* 2006;21:975–8.
- 194 Estacio RO, Jeffers BW, Gifford N, Schrier RW. Effect of blood pressure control on diabetic microvascular complications in patients with hypertension and type 2 diabetes. *Diabetes Care* 2000;23(Suppl 2):B54–64.
- 195 Wright JT Jr, Bakris G, Greene T *et al.* Effect of blood pressure lowering and antihypertensive drug class on progression of hypertensive kidney disease: results from the AASK trial. *JAMA* 2002;288:2421–31.
- 196 Ismail-Beigi F, Craven TE, O'Connor PJ *et al.* Combined intensive blood pressure and glycemic control does not produce an additive benefit on microvascular outcomes in type 2 diabetic patients. *Kidney Int* 2012;81:586–94.
- 197 Brouwer TF, Vehmeijer JT, Kalkman DN, Berger WR, van den Born BH, Peters RJ, Knops RE. Intensive blood pressure lowering in patients with and patients without type 2 diabetes: A pooled analysis from two randomized trials. *Diabetes Care* 2018;41:1142–8.
- 198 Basu S, Sussman JB, Rigdon J, Steimle L, Denton BT, Hayward RA. Benefit and harm of intensive blood pressure treatment: Derivation and validation of risk models using data from the SPRINT and ACCORD trials. *PLoS Med* 2017;14: e1002410.
- 199 Buyadaa O, Magliano DJ, Salim A, Koye DN, Shaw JE. Risk of rapid kidney function decline, all-cause mortality, and major cardiovascular events in nonalbuminuric chronic kidney disease in type 2 diabetes. *Diabetes Care* 2020;43:122–9.
- 200 de Zeeuw D, Remuzzi G, Parving HH *et al.* Albuminuria, a therapeutic target for cardiovascular protection in type 2 diabetic patients with nephropathy. *Circulation* 2004;110:921–7.
- 201 Bakris GL, Weir MR, Shanifar S *et al.* Effects of blood pressure level on progression of diabetic nephropathy: results from the RENAAL study. *Arch Intern Med* 2003;163:1555–65.
- 202 Holtkamp FA, de Zeeuw D, de Graeff PA *et al.* Albuminuria and blood pressure, independent targets for cardioprotective therapy in patients with diabetes and nephropathy: a post hoc analysis of the combined RENAAL and IDNT trials. *Eur Heart J* 2011;32:1493–9.
- 203 Berl T, Hunsicker LG, Lewis JB *et al.* Impact of achieved blood pressure on cardiovascular outcomes in the Irbesartan Diabetic Nephropathy Trial. *J Am Soc Nephrol* 2005;16:2170–9.
- 204 Gaede P, Vedel P, Larsen N *et al.* Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003;348:383–93.
- 205 Wheeler DC, Becker GJ. Summary of KDIGO guideline. What do we really know about management of blood pressure in patients with chronic kidney disease? *Kidney Int* 2013;83:377–83.
- 206 Estacio RO, Jeffers BW, Hiatt WR *et al.* The effect of nisoldipine as compared with enalapril on cardiovascular outcomes in patients with non-insulin-dependent diabetes and hypertension. *N Engl J Med* 1998;338:645–52.

- 207 Xie X, Liu Y, Perkovic V *et al.* Renin-Angiotensin System Inhibitors and Kidney and Cardiovascular Outcomes in Patients With CKD: A Bayesian Network Meta-analysis of Randomized Clinical Trials. *Am J Kidney Dis* 2016;67:728–41.
- 208 Cheng J, Zhang W, Zhang X *et al.* Effect of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers on all-cause mortality, cardiovascular deaths, and cardiovascular events in patients with diabetes mellitus: a meta-analysis. *JAMA Intern Med* 2014;174:773–85.
- 209 Imai E, Chan JC, Ito S *et al.* Effects of olmesartan on renal and cardiovascular outcomes in type 2 diabetes with overt nephropathy: a multicentre, randomised, placebo-controlled study. *Diabetologia* 2011;54:2978–86.
- 210 Palmer SC, Mavridis D, Navarese E *et al.* Comparative efficacy and safety of blood pressure-lowering agents in adults with diabetes and kidney disease: a network meta-analysis. *Lancet* 2015;385:2047–56.
- 211 Effects of ramipril on cardiovascular and microvascular outcomes in people with diabetes mellitus: Results of the HOPE study and MICRO-HOPE substudy. heart outcomes prevention evaluation study investigators. *Lancet* 2000;355:253–9.
- 212 Qiao Y, Shin JJ, Chen TK, Inker LA, Coresh J, Alexander GC, Jackson JW, Chang AR, Grams ME. Association between renin-angiotensin system blockade discontinuation and all-cause mortality among persons with low estimated glomerular filtration rate. *JAMA Intern Med* 2020.
- 213 Bhandari S, Ives N, Brettell EA, Valente M, Cockwell P, Topham PS, Cleland JG, Khwaja A, El Nahas M. Multicentre randomized controlled trial of angiotensin-converting enzyme inhibitor/angiotensin receptor blocker withdrawal in advanced renal disease: The STOP-ACEi trial. *Nephrol Dial Transplant* 2016;31:255–261.
- 214 van den Meiracker AH, Baggen RG, Pauli S *et al.* Spironolactone in type 2 diabetic nephropathy: Effects on proteinuria, blood pressure and renal function. *J Hypertens* 2006;24:2285–92.
- 215 Bushinsky DA, Williams GH, Pitt B *et al.* Patiromer induces rapid and sustained potassium lowering in patients with chronic kidney disease and hyperkalemia. *Kidney Int* 2015;88:1427–33.
- 216 Bakris GL, Pitt B, Weir MR *et al.* Effect of Patiromer on Serum Potassium Level in Patients With Hyperkalemia and Diabetic Kidney Disease: The AMETHYST-DN Randomized Clinical Trial. *JAMA* 2015;314:151–61.
- 217 Weir MR, Bakris GL, Bushinsky DA *et al.* Patiromer in patients with kidney disease and hyperkalemia receiving RAAS inhibitors. *N Engl J Med* 2015;372:211–21.
- 218 Pitt B, Bakris GL, Bushinsky DA *et al.* Effect of patiromer on reducing serum potassium and preventing recurrent hyperkalaemia in patients with heart failure and chronic kidney disease on RAAS inhibitors. *Eur J Heart Fail* 2015;17:1057–65.
- 219 Weir MR, Bakris GL, Gross C *et al.* Treatment with patiromer decreases aldosterone in patients with chronic kidney disease and hyperkalemia on renin-angiotensin system inhibitors. *Kidney Int* 2016;90:696–704.
- 220 Agarwal R, Rossignol P, Romero A *et al.* Patiromer versus placebo to enable spironolactone use in patients with resistant hypertension and chronic kidney disease (AMBER): a phase 2, randomised, double-blind, placebo-controlled trial *Lancet* 2019;394:1540–50.
- 221 Ganesan C, Pao AC. Spironolactone plus patiromer: proceed with caution. *Lancet*. 2019 Oct 26;394(10208):1486–1488. doi: 10.1016/S0140-6736(19)32485-7. PMID: 31657725.

- 222 McMahon EJ, Bauer JD, Hawley CM *et al.* A randomized trial of dietary sodium restriction in CKD. *J Am Soc Nephrol* 2013;24:2096–2103.
- 223 de Brito-Ashurst I, Perry L, Sanders TA *et al.* The role of salt intake and salt sensitivity in the management of hypertension in South Asian people with chronic kidney disease: a randomised controlled trial. *Heart* 2013;99:1256–60.
- 224 Van Huffel L, Tomson CR, Ruige J, Nistor I, Van Biesen W, Bolignano D. Dietary restriction and exercise for diabetic patients with chronic kidney disease: A systematic review. *PLoS One* 2014;9: e113667.
- 225 MacMahon S, Peto R, Cutler J *et al.* Blood pressure, stroke, and coronary heart disease. Part 1, Prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet* 1990;335:765–74.
- 226 Kannel WB. Blood pressure as a cardiovascular risk factor: prevention and treatment. *JAMA* 1996;275:1571–6.
- 227 Kannel WB, McGee DL. Diabetes and cardiovascular risk factors: the Framingham study. *Circulation* 1979;59:8–13.
- 228 Foley RN, Parfrey PS, Sarnak MJ. Clinical epidemiology of cardiovascular disease in chronic renal disease. *Am J Kidney Dis* 1998;32(Suppl 3):S112–9.
- 229 Foley RN, Murray AM, Li S *et al.* Chronic kidney disease and the risk for cardiovascular disease, renal replacement, and death in the United States Medicare population, 1998 to 1999. *J Am Soc Nephrol* 2005;16:489–95.
- 230 M Kessler, F Zannad, P Leheret *et al.* Predictors of cardiovascular events in patients with end-stage renal disease: an analysis from the Fosinopril in Dialysis study. *Nephrol Dial Transplant* 2007;22:3573–79.
- 231 Stel VS, van de Luijngaarden MW, Wanner C *et al.* The 2008 ERA-EDTA Registry Annual Report – a précis. *NDT Plus* 2011;4:1–13.
- 232 Steenkamp R, Rao A, Fraser S. UK Renal Registry 18th Annual Report, Chapter 5: Survival and Causes of Death in UK Adult Patients on Renal Replacement Therapy in 2014. *Nephron* 2016;132(Suppl 1):145–54.
- 233 Vonesh EF, Snyder JJ, Foley RN *et al.* The differential impact of risk factors on mortality in hemodialysis and peritoneal dialysis. *Kidney Int* 2004;66:2389–401.
- 234 Liem YS, Wong JB, Hunink MG *et al.* Comparison of hemodialysis and peritoneal dialysis survival in the Netherlands. *Kidney Int* 2007;71:153–8.
- 235 Collins AJ. Cardiovascular mortality in end-stage renal disease. *Am J Med Sci* 2003;325:163–7.
- 236 Sarnak MJ. Cardiovascular complications in chronic kidney disease. *Am J Kidney Dis* 2003;41(Suppl):11–7.
- 237 Collins AJ, Foley R, Herzog C *et al.* Excerpts from the United States Renal Data System 2007 annual data report. *Am J Kidney Dis* 2008;51(Suppl 1):S1–320.
- 238 Agarwal R. Hypertension and survival in chronic hemodialysis patients – past lessons and future opportunities. *Kidney Int* 2005;67:1–13.
- 239 Hirakata H, Onoyama K, Hori K *et al.* The hemodynamic and humoral responses to tilting in diabetic patients on chronic hemodialysis treatment. *Clin Nephrol* 1987;27:298–303.
- 240 Itsuko I, Hirakata H, Sugimori H *et al.* Hemodialysis causes severe orthostatic reduction in cerebral blood flow velocity in diabetic patients. *Am J Kidney Dis* 1999;34:1096–104.
- 241 Peixoto AJ, Santos SF, Mendes RB *et al.* Reproducibility of ambulatory blood pressure monitoring in hemodialysis patients. *Am J Kidney Dis* 2000;36:983–90.

- 242 Dolan E, Stanton A, Thijs L *et al.* Superiority of ambulatory over clinic blood pressure measurement in predicting mortality: the Dublin outcome study. *Hypertension* 2005;46:156–61.
- 243 Agarwal R, Peixoto AJ, Santos SF, Zoccali C. Pre- and postdialysis blood pressures are imprecise estimates of interdialytic ambulatory blood pressure. *Clin J Am Soc Nephrol* 2006;1:389–98.
- 244 Agarwal R, Peixoto AJ, Santos SF, Zoccali C. Out-of-office blood pressure monitoring in chronic kidney disease. *Blood Press Monit* 2009;14:2–11.
- 245 Moriya H, Ohtake T, Kobayashi S. Aortic stiffness, left ventricular hypertrophy and weekly averaged blood pressure (WAB) in patients on haemodialysis. *Nephrol Dial Transplant* 2007;22:1198–204.
- 246 Rahman M, Griffin V, Heyka R, Hoit B. Diurnal variation of blood pressure; reproducibility and association with left ventricular hypertrophy in hemodialysis patients. *Blood Press Monit* 2005;10:25–32.
- 247 Zoccali C, Mallamaci F, Tripepi G *et al.* Prediction of left ventricular geometry by clinic, pre-dialysis and 24-h ambulatory BP monitoring in hemodialysis patients: CREED investigators. *J Hypertens* 1999;17:1751–8.
- 248 Agarwal R, Brim NJ, Mathenthiran J *et al.* Out-of-haemodialysis-unit blood pressure is a superior determinant of left ventricular hypertrophy. *Hypertension* 2006;47:62–8.
- 249 Amar J, Vernier I, Rossignol E *et al.* Nocturnal blood pressure and 24-hour pulse pressure are potent indicators of mortality in hemodialysis patients. *Kidney Int* 2000;57:2485–91.
- 250 Alborzi P, Patel N, Agarwal R. Home blood pressures are of greater prognostic value than hemodialysis unit recordings. *Clin J Am Soc Nephrol* 2007;2:1228–34.
- 251 Verdecchia P, Angeli F. How can we use the results of ambulatory blood pressure monitoring in clinical practice? *Hypertension* 2005;46:25–6.
- 252 Liu M, Takahashi H, Morita Y *et al.* Non-dipping is a potent predictor of cardiovascular mortality and is associated with autonomic dysfunction in haemodialysis patients. *Nephrol Dial Transplant* 2003;18:563–9.
- 253 Port FK, Hulbert-Shearon TE, Wolfe RA *et al.* Predialysis blood pressure and mortality risk in a national sample of maintenance hemodialysis patients. *Am J Kidney Dis* 1999;33:507–17.
- 254 Goldfarb-Rumyantzev AS, Baird BC, Leypoldt JK, Cheung AK. The association between BP and mortality in patients on chronic peritoneal dialysis. *Nephrol Dial Transplant* 2005;20:1693–1701.
- 255 Kalantar-Zadeh K, Kilpatrick RD, Kopple JD. Reverse epidemiology of blood pressure in dialysis patients. *Kidney Int* 2005;67:2067; author reply 2067–8.
- 256 Li Z, Lacson E Jr, Lowrie EG *et al.* The epidemiology of systolic blood pressure and death risk in hemodialysis patients. *Am J Kidney Dis* 2006;48:606–15.
- 257 Robinson BM, Tong L, Zhang J, Wolfe RA, Goodkin DA, Greenwood RN, Kerr PG, Morgenstern H, Li Y, Pisoni RL, Saran R, Tentori F, Akizawa T, Fukuhara S, Port FK. Blood pressure levels and mortality risk among hemodialysis patients in the Dialysis Outcomes and Practice Patterns Study. *Kidney Int* 2012;82(5):570–80.
- 258 Bansal N, McCulloch CE, Lin F, Alper A, Anderson AH, Cuevas M, Go AS, Kallem R, Kusek JW, Lora CM, Lustigova E, Ojo A, Rahman M, Robinson-Cohen C, Townsend RR, Wright J, Xie D, Hsu CY. Blood Pressure and Risk of Cardiovascular Events in Patients on Chronic Hemodialysis: The CRIC Study (Chronic Renal Insufficiency Cohort). *Hypertension* 2017;70(2):435–443.

- 259 Charra B, Calemard E, Ruffet M *et al.* Survival as an index of adequacy of dialysis. *Kidney Int* 1992;41:1286–91.
- 260 Suzuki H, Kanno Y, Sugahara S *et al.* Effect of angiotensin receptor blockers on cardiovascular events in patients undergoing hemodialysis: an open-label randomized controlled trial. *Am J Kidney Dis* 2008;52:501–6.
- 261 Zannad F, Kessler M, Leheret P *et al.* Prevention of cardiovascular events in end-stage renal disease: results of a randomized trial of fosinopril and implications for future studies. *Kidney Int* 2006;70:1318–24.
- 262 Cice G, Ferrara L, D’Andrea A *et al.* Carvedilol increases two-year survival in dialysis patients with dilated cardiomyopathy: a prospective, placebo-controlled trial. *J Am Coll Cardiol* 2003;41:1438–44.
- 263 Tepel M, Hopfenmueller W, Scholze A *et al.* Effect of amlodipine on cardiovascular events in hypertensive haemodialysis patients. *Nephrol Dial Transplant* 2008;23:3605–12.
- 264 Heerspink HJ, Ninomiya T, Zoungas S *et al.* Effect of lowering blood pressure on cardiovascular events and mortality in patients on dialysis: a systematic review and meta-analysis of randomised controlled trials. *Lancet* 2009;373:1009–15.
- 265 Agarwal R, Sinha AD. Cardiovascular protection with antihypertensive drugs in dialysis patients: systematic review and meta-analysis. *Hypertension* 2009;53:860–6.
- 266 Agarwal R, Sinha AD, Pappas MK *et al.* Hypertension in hemodialysis patients treated with atenolol or lisinopril: a randomized controlled trial. *Nephrol Dial Transplant* 2014;29:672–81.
- 267 Levin NW, Kotanko P, Eckardt KU *et al.* Blood pressure in chronic kidney disease stage 5D – report from a Kidney Disease: Improving Global Outcomes controversies conference. *Kidney Int* 2007;77:273–84.
- 268 K/DOQI Workgroup. K/DOQI clinical practice guidelines for cardiovascular disease in dialysis patients. *Am J Kidney Dis* 2005;45(Suppl 3):S1–153.
- 269 Klassen PS, Lowrie EG, Reddan DN *et al.* Association between pulse pressure and mortality in patients undergoing maintenance haemodialysis. *JAMA* 2002;287:1548–55.
- 270 Wang, AY, Brimble, KS, Brunier, G, Holt, SG, Jha, V, Johnson, DW, Kang, SW, Kooman, JP, Lambie, M, McIntyre, C, Mehrotra, R, Pecoits-Filho, R. ISPD Cardiovascular and Metabolic Guidelines in Adult Peritoneal Dialysis Patients Part I - Assessment and Management of Various Cardiovascular Risk Factors. *Perit Dial Int* 2015;35(4):379-87.
- 271 Wilson J, Shah T, Nissenson AR. Role of sodium and volume in the pathogenesis of hypertension in hemodialysis. *Seminars in dialysis* 2004;17:260–4.
- 272 Agarwal R, Weir MR. Dry-Weight: A Concept Revisited in an Effort to Avoid Medication-Directed Approaches for Blood Pressure Control in Hemodialysis Patients. *CJASN* 2010;5:1255–60.
- 273 Thomson GE, Waterhouse K, McDonald HP Jnr, Friedman EA. Hemodialysis for chronic renal failure. Clinical observations. *Arch Intern Med* 1967;120:153–67.
- 274 Charra B, Bergström J, Scribner BH. Blood pressure control in dialysis patients: importance of the lag phenomenon. *Am J Kidney Dis* 1998;32:720–4.
- 275 Ozkahya M, Toz H, Qzerkan F *et al.* Impact of volume control on left ventricular hypertrophy in dialysis patients. *J Nephrology* 2002;15:655–60.
- 276 Aşci G, Özkahya M, Duman S *et al.* Volume control associated with better cardiac function in long-term peritoneal dialysis patients. *Peritoneal dialysis International* 2006;26: 85–8.

- 277 Agarwal R, Alborzi P, Satyan S, Light RP. Dry-weight reduction in hypertensive haemodialysis patients (DRIP): a randomized, controlled trial. *Hypertension* 2009;53:500–7.
- 278 Luik AJ, Charra B, Katzarski K *et al.* Blood pressure control and hemodynamic changes in patients on long time dialysis treatment. *Blood Purif* 1998;16:197–209.
- 279 Zilch O, Vos PF, Oey PL *et al.* Sympathetic hyperactivity in haemodialysis patients is reduced by short daily haemodialysis. *J Hypertens* 2007;25:1285–9.
- 280 Culeton BF, Walsh M, Klarenbach SW *et al.* Effect of frequent nocturnal hemodialysis vs conventional hemodialysis on left ventricular mass and quality of life: a randomized controlled trial. *JAMA* 2007;298:1291–9.
- 281 Pierratos A, Ouwendyk M, Francoeur R *et al.* Nocturnal hemodialysis: three-year experience. *J Am Soc Nephrol* 1998;9:859–68.
- 282 Shoji T, Tsubakihara Y, Masamitsu F *et al.* Hemodialysis-associated hypotension as an independent risk factor for two-year mortality in hemodialysis patients. *Kidney Int* 2004;66:1212–20.
- 283 Bragg-Gresham JL, Fissell RB, Mason NA *et al.* Diuretic use, residual renal function, and mortality among hemodialysis patients in the Dialysis Outcomes and Practice Pattern Study (DOPPS). *Am J Kidney Dis* 2007;49:426–31.
- 284 Movilli E, Gaggia P, Zubani R *et al.* Association between high ultrafiltration rates and mortality in uraemic patients on regular haemodialysis. A 5-year prospective observational multicentre study. *Nephrol Dial Transplant* 2007;22:3547–52.
- 285 Onofriescu M, Hogas S, Voroneanu L *et al.* Bioimpedance-guided fluid management in maintenance hemodialysis: a pilot randomized controlled trial. *Am J Kidney Dis* 2014;64:111–8.
- 286 Roueff S, Martin E, Chauffert ML *et al.* Brain natriuretic peptide variations are linked to volume status in hemodialysis patients. *Clin Nephrol* 2008;70:508–13.
- 287 Brennan JM, Ronan A, Goonewardena S *et al.* Handcarried ultrasound measurement of the inferior vena cava for assessment of intravascular volume status in the outpatient hemodialysis clinic. *Clin J Am Soc Nephrol* 2006;1:749–53.
- 288 Picano E, Frassi F, Agricola E *et al.* Ultrasound lung comets: a clinically useful sign of extravascular lung water. *J Am Soc Echocardiogr* 2006;19(3):356–63.
- 289 Davies SJ, Davenport A. The role of bioimpedance and biomarkers in helping to aid clinical decision-making of volume assessments in dialysis patients. *Kidney Int* 2014;86(3):489–96.
- 290 Hegstrom RM, Murray JS, Pendras JP *et al.* Hemodialysis in the treatment of chronic uremia. *Trans Am Soc Artif Intern Organs* 1961;7:136–52.
- 291 Rigby-Matthews A, Scriber BH, Ahmad S. Control of interdialytic weight gain does not require fluid restriction in hemodialysis (HD) patients (Abstract). *J Am Soc Nephrol* 1999;10:267A.
- 292 Craswell PW, Hird VM, Judd PA *et al.* Plasma renin activity and blood pressure in 89 patients receiving maintenance haemodialysis therapy. *Br Med J* 1972;4:749–53.
- 293 Covic A, Goldsmith DJ, Venning MC, Ackrill P. Long-hours home haemodialysis – the best renal replacement therapy method? *QJM* 1999;92:251–260.
- 294 Ozkahya M, Ok E, Toz H *et al.* Long-term survival rates in haemodialysis patients treated with strict volume control. *Nephrol Dial Transplant* 2006;21:3506–13.

- 295 Kayikcioglu M, Tumuklu M, Ozkahya M *et al*. The benefit of salt restriction in the treatment of end-stage renal disease by haemodialysis. *Nephrol Dial Transplant* 2009;24:956–62.
- 296 Maduell F, Navarro V. Dietary salt intake and blood pressure control in haemodialysis patients. *Nephrol Dial Transplant* 2000;15:2063.
- 297 Mc Causland FR, Waikar SS, Brunelli SM. Increased dietary sodium is independently associated with greater mortality among prevalent hemodialysis patients. *Kidney Int* 2012;82:204–11.
- 298 Medcalf JF, Harris KP, Walls J. Role of diuretics in the preservation of residual renal function in patients on continuous ambulatory peritoneal dialysis. *Kidney Int* 2001;59:1128–33.
- 299 Ni X, Zhang J, Zhang P *et al*. Effects of spironolactone on dialysis patients with refractory hypertension: a randomized controlled study. *J Clin Hypertension (Greenwich)* 2014;16:658–63.
- 300 Georgianos PI, Agarwal R. Pharmacotherapy of hypertension in chronic dialysis patients. *Clin J Am Soc Nephrol* 2016;11:2062–27.
- 301 University Hospital Brest, Clinical Trial: NCT01848639: ALdosterone Antagonist Chronic HEModialysis Interventional Survival Trial (ALCHEMIST), 2013. Available from <https://clinicaltrials.gov/ct2/show/NCT01848639> [Accessed February 2017].